

# THE PHYSICIAN'S GUIDE TO



## Treating Enuresis and Encopresis with the Modified O'Regan Protocol

**"M.O.P. works radically better than anything else."**

– James Sander, M.D., Director of Pediatric Urology,  
Doctors Hospital at Renaissance, Edinburg, TX


**"Families and kids are  
a lot more receptive to  
M.O.P. than I would  
have thought."**

– Irina Stanasel, M.D., pediatric  
urologist, UT Southwestern  
Medical Center, Dallas, TX



By Steve Hodges, M.D.

Associate Professor of Pediatric Urology  
Wake Forest University School of Medicine



**"It is my mission to get  
the word out about  
how incredibly effective  
M.O.P. is."**

– Erin Wetjen, PT, specialist  
in pediatric incontinence,  
Mayo Clinic, Rochester, MN

**"No enuresis treatment works as  
well as M.O.P. — not even close."**

– Victoriano Romero, M.D., urologist,  
Redding Urologic Associates, Redding, CA

# Contents



A Letter to Colleagues	1
The Modified O'Regan Protocol in Brief	2
• The premise behind M.O.P.	2
• M.O.P. guidelines and safety considerations	3
• Origins of M.O.P.	4
• What is "modified" about M.O.P.	5
Five Lessons from a Decade Prescribing M.O.P.	6
• Rectal diameter is critical	6
• Enemas are far more effective than laxatives	7
• Daily enemas do not cause dependence or electrolyte imbalance	8
• Enemas are not "abusive" or "traumatic"	9
• Alternatives to Miralax can work just as well	10
The M.O.P. Progression Chart	11
Resources for Parents	12
Full Text of Key Studies	18

September 2020

**Department of Urology**

Medical Center Boulevard  
Winston-Salem, NC 27157  
p 336.716.0662  
f 336.716.5711  
www.WakeHealth.edu/urology

Dear Colleague,

I am writing to share a treatment protocol for enuresis and encopresis that works far better for my patients than Miralax, medication, alarms, or other treatments I was taught in medical school. It is an enema-based regimen called M.O.P., short for the Modified O'Regan Protocol. M.O.P. is based on the research of pediatric nephrologist Dr. Sean O'Regan, who practiced at the University of Montreal Hôpital Sainte-Justine in the 1980s and is now retired.

Dr. O'Regan's original and highly successful protocol lasted 90 days:

- 30 consecutive days of enemas
- 30 days of enemas every other day
- 30 days of enemas twice a week

I have "modified" Dr. O'Regan's protocol based on my own research and clinical experience, as well as discussions with Dr. O'Regan himself. Most significantly, I recommend:

- adding a daily osmotic laxative to the enema regimen (except for the first 2, in the case of encopresis, as laxatives often make matters worse).
- extending the daily enema phase if needed, tapering only after the child has achieved 7 dry days/nights *and* has completed 30 enemas.
- shifting to large-volume enemas if no progress after 30 days.
- adding daily Ex-Lax during the tapering phase or earlier if the child is not spontaneously pooping daily.

Physicians unfamiliar with this protocol may consider it overly aggressive, perhaps even traumatic for children. Some worry daily enemas will cause dependency and/or electrolyte imbalance. But published studies and my experience show these concerns are unwarranted. I have treated thousands of children with M.O.P., without incident and with excellent success.

Enuresis patients are often told, "Don't worry, you'll outgrow it," but my patient load includes countless tweens and teens whose accidents were dismissed for years. In fact, I have so many teenage patients that I recently published a book just for them: *M.O.P. for Teens and Tweens*. Though we, as physicians, often consider enuresis and encopresis to be relatively minor issues, these conditions are deeply distressing and embarrassing for children and their parents. I have made enuresis and encopresis the focus of my research and clinical practice and am eager to share with colleagues the treatments, however unconventional, that have worked so well with my patients.

This packet briefly explains the rationale and scientific support for M.O.P. The regimen is not so much a protocol as a process, with many permutations, described more thoroughly in *The M.O.P. Book: Anthology Edition*, a guide for parents. I urge physicians to become familiar with all the variations. I am happy to email you a PDF of *The M.O.P. Book* and *M.O.P. for Teens and Tweens*. — just ask!

I'm always available to discuss enema safety and effectiveness with colleagues. Email me and we'll set up a time to talk!



Steve Hodges, M.D.  
shodges@wakehealth.edu

**Anthony Atala MD**  
W H Boyce Professor and Chair  
**Gopal H. Badlani MD**  
Professor  
Vice Chair, Clinical Affairs  
**K. C. Balaji MD**  
Professor  
Director, V.A. Urology Service  
**Ronald L. Davis, III MD, MBA**  
Associate Professor  
**Raymond B. Dyer MD**  
UroRadiology  
**Robert J. Evans MD**  
Associate Professor, Clinic Director  
**Barry I. Freedman MD**  
Nephrology  
**Jorge Gutierrez MD**  
Professor  
Director, Endourology  
**Ashok K. Hemal MD**  
Professor  
Director, Robotics  
**Steve J. Hodges MD**  
Associate Professor  
**Stuart S. Howards MD**  
Professor  
**Stanley J. Kogan MD**  
Professor  
**Samuel Lentz MD**  
UroGynecology  
**Catherine Matthews MD**  
Professor  
Co-Director, Female Pelvic Health  
**John D. McConnell MD**  
Professor and CEO  
Wake Forest Baptist Medical Center  
**Majid Mirzazadeh MD**  
Assistant Professor  
**Robert G. Moore MD**  
Clinical Associate Professor  
**Robert A. Older MD**  
Professor  
**Michael A. Olympio MD**  
UroAnesthesia  
**Candace Parker-Autry MD**  
UroGynecology  
**Daniel B. Rukstalis MD**  
Professor  
Director, Residency Program  
**Hooman Sadri-Ardekani MD, PhD**  
Instructor  
**Robert Stratta MD**  
Transplant Surgery  
**Allston J. Stubbs MD**  
Clinical Associate Professor  
**Ryan Terlecki MD**  
Associate Professor  
**Christopher Y. Thomas MD**  
GU Oncology

**Research Faculty**  
Karl-Erik Andersson MD, PhD  
Colin E. Bishop PhD  
Yong Chen PhD  
Steven Kridel PhD  
George Kulik MD, DVM  
Guangchao Sui PhD  
James J. Yoo MD, PhD  
Stephen J. Walker, PhD  
J. Kouly Williams DVM  
Weixin Zhao, MD

# The Modified O'Regan Protocol in Brief

## The premise behind M.O.P.

The Modified O'Regan Protocol is based on a simple, proven premise: **The cause of all encopresis and virtually all enuresis — primary and secondary, nocturnal and daytime — is chronic constipation. The distinctions among these forms of enuresis are meaningless.**


Despite what the AAP and other organizations suggest, **it is not “deep sleep,” an underdeveloped bladder, excess urine production, behavioral issues, stress, or heredity that is the root cause of these conditions.** (While it is true that bedwetting can run in families, it is the tendency toward constipation and the sensitivity of the bladder to a stretched rectum — not enuresis itself — that may be passed on.) Below, I cite the research demonstrating constipation is almost always the culprit. Research does not support any other hypothesis, and to suggest the causes are unknown, as some organizations do, ignores definitive research to the contrary. (In rare cases, enuresis is caused by polyuria from diabetes or neurological/anatomical abnormalities such as spina bifida, tethered cord, or imperforate anus, but those cases should be obvious.)

**Though nocturnal enuresis is common, it is absolutely not “normal” and must be treated early and aggressively.** I recommend treating nocturnal enuresis beginning at age 4. Any child who struggles with toilet training or has daytime accidents after toilet training must also be treated aggressively, even at age 2 or 3. The holding habit is deeply ingrained, and left untreated, many children will experience worsening symptoms over time. Children with both daytime and nighttime symptoms are the least likely to outgrow bedwetting.<sup>1</sup> However, you cannot assume that any given child, even one with only nocturnal enuresis, will outgrow the condition. My clinic and Facebook support groups are filled with families who were assured their child would stop wetting eventually. But that day never came.

### How constipation causes enuresis:

An abnormally dilated rectum in susceptible children causes uninhibited bladder contractions. I order KUB X-rays for all my enuresis patients to prove this point to parents and to establish a baseline rectal diameter measurement. A diameter above 3 cm signals constipation. Most of my patients measure at least 6 cm.

Though numerous articles distinguish between primary and secondary enuresis, this distinction is needless, as the cause is the same, which X-rays plainly show.



1 Yeung, C., Sreedhar, B., Sihoe, J., Sit, F., & Lam, J. (2006, April 6). Differences in characteristics of nocturnal enuresis between children and adolescents: a critical appraisal from a large epidemiological study. *BJU International*, 97(5), 1069-1073. doi: <https://doi.org/10.1111/j.1464-410X.2006.06074.x>

**How constipation causes encopresis:** The stool-stuffed rectum becomes chronically stretched, to the point of losing tone and sensation. These kids never fully empty, nor do they feel the urge to poop. Referrals to behavioral therapists are unwarranted and only serve to frustrate families.

The way to resolve both enuresis and encopresis is to clean out the rectum on a daily basis for several weeks, even months, allowing the rectum to remain clear, shrink back to size, stop aggravating the bladder, and regain tone and sensation.

**The “first-line” therapies for enuresis – alarms and medication – do not achieve these goals.**

And treating chronic constipation with osmotic laxatives alone is typically insufficient for children with enuresis and encopresis. Even when it does work, the results tend to be fleeting. In my experience, physicians typically underestimate the severity of the constipation in enuresis and encopresis patients and therefore undertreat these patients, as I did for many years.

In well over a decade of treating these conditions, I have found that the Modified O’Regan Protocol is by far the most effective remedy for enuresis and encopresis, as well as recurrent urinary tract infections. Many of my patients have taken high doses of Miralax for years and years, only to have their enuresis resolve within a couple of months on **M.O.P.** Encopresis, often made worse by Miralax, typically resolves within a month on **M.O.P.**, though the tapering phase is important.

## **M.O.P. guidelines and safety considerations**

Following are the key guidelines for the Modified O’Regan Protocol.

### **1.) Do daily enemas for at least 30 days.**

Taper only when the child has completed 30 days of enemas AND remains accident-free for at least 7 consecutive days. If the child makes no progress in the first month, switch to liquid glycerin suppositories or large-volume enemas with glycerin and/or Castile soap (**M.O.P.+**). Optional but helpful: periodic overnight oil-retention enemas with olive oil or mineral oil, followed by a morning stimulant enema.

### **2.) After at least 30 days of enemas AND 7 consecutive days and nights of dryness, taper to one enema every other day for another 30 days.**

If accidents recur, resume daily enemas until achieving 7 consecutive accident-free days.

### **3.) After a second 30 days of dryness, taper to enemas twice a week for 30 days before stopping.**

If accidents recur, resume daily enemas until achieving 7 consecutive accident-free days, and then taper to every other day for 30 days.

### **4.) Take an osmotic laxative daily to keep stool soft.**

Use PEG 3350, lactulose, magnesium citrate, magnesium hydroxide, or other laxative of choice. Take a full dose daily or half a dose twice a day. Continue daily for 6 months after **M.O.P.** For encopresis patients, hold off on an osmotic laxative for the first 2 weeks.

### **Important safety considerations:**

- 1.) Enemas should not be performed on a child with kidney disease.
- 2.) Children with other chronic diseases must consult with a physician before doing enemas.
- 3.) Children should not be given more than one enema per day. The exception: Overnight oil enemas. These do not “count” as a second enema, as they only soften stool rather than stimulate a bowel movement.

## Origins of M.O.P.

**M.O.P.** is named for Dr. Sean O'Regan, an Irish pediatric nephrologist who practiced in Montreal in the 1980s. It was Dr. O'Regan who first proved constipation is the cause of enuresis, encopresis, and recurrent UTIs and who pioneered the use of daily enemas to resolve these conditions.

Dr. O'Regan did not set out to show constipation causes enuresis. He was simply trying to figure out why his 5-year-old was wetting the bed nightly and how to resolve the problem. At this time, children who wet the bed were thought to have psychological and/or anatomical problems, such as an excessively narrow bladder neck. Dr. O'Regan did not accept either explanation and began searching for answers at the McGill University Medical Library. He was surprised to find several references, dating back to the 1890s, to a connection between constipation and urinary problems.

Dr. O'Regan felt he was onto something. So he had a colleague, Dr. Salam Yazbeck, test his son's rectum using anorectal manometry, the most reliable indicator of constipation. Dr. Yazbeck reported to Dr. O'Regan, "The kid's got no rectal tone." Dr. O'Regan began giving his son nightly enemas. Within a week, the boy was having his first dry nights. Within 2 months, he'd stopped wetting completely.

Based on this success, Drs. O'Regan and Yazbeck began a series of studies at the University of Montreal using the protocol he tried with his son: daily enemas for a month, followed by a month of enemas every other day, followed by a third month of twice-weekly enemas. They got the word out to local pediatricians and attracted virtually the entire French Canadian population of children with urinary problems. Advancing the 1960s research, their studies showed constipation was the cause of urinary problems in children and that treating the constipation with enemas resolved these problems in dramatic fashion.

**I urge you to read the full text of Dr. O'Regan's studies, posted at [BedwettingAndAccidents.com](http://BedwettingAndAccidents.com).**

In one investigation, published in 1985, Dr. O'Regan tracked 47 girls, average age 8, with recurrent UTIs who were shown by anorectal manometry to be severely constipated.<sup>2</sup> In addition, most of these girls had encopresis and/or enuresis. Dr. O'Regan prescribed the enema regimen he had used with his son. Within 3 months, 44 of the 47 girls no longer were having UTIs. Encopresis resolved in 20 of the 21 patients with this condition, and 22 of the 32 girls with enuresis stopped wetting. Among the girls who didn't improve, most of their parents admitted to not following the enema regimen fully.

**Dr. O'Regan got no pushback from physicians regarding the use of enemas.** None suggested enemas would damage a child's physical or emotional health or that oral laxatives would be safer or more effective. (Though Miralax wasn't available back then, senna, magnesium, and castor oil were among common oral remedies.) "We knew the root cause of bedwetting was incomplete rectal emptying," Dr. O'Regan told me in an interview, "and enemas were the only way to solve the problem."

Over the years, I have come to agree. I've been so fascinated by the modern-day medical community's alarm about enemas that I recently called Dr. O'Regan, now retired in Arizona, and asked him if any of his patients ever suffered complications on the regimen. He told me, "Our only complication was a 7-year-old girl who clogged the toilet at our hospital after an enema. She was legendary."

The damage was to the plumbing, not the patient!

2 O'Regan, S., Yazbeck, S., & Schick, E. (1985). Constipation, bladder instability, urinary tract infection syndrome. *Clinical nephrology*, 23(3), 152–154. <https://pubmed.ncbi.nlm.nih.gov/3987104>.

## What is “modified” about **M.O.P.**

Dr. O'Regan had remarkable success with his 90-day, step-down regimen. For years I had success, too, but less than what Dr. O'Regan reported. In examining my patient load, I suspected this was because childhood constipation is more severe and more prevalent today than it was in Dr. O'Regan's time.

Childhood obesity rates have tripled since the 1980s, due to our kids' highly processed diet and low activity levels — two factors also driving the constipation epidemic. In addition, more children today attend preschool, and most facilities require toilet training by age 3, prompting parents to train before the child is ready. My own research shows children trained before age 2 have triple the risk of developing chronic constipation and enuresis.<sup>3</sup>

Finally, public school bathrooms today are scarier and filthier than they were in the 1980s, and school restroom policies are more restrictive, so fewer school-age kids are using the toilet at school. In a UCSF survey of more than 4,000 elementary teachers, 88% said they encourage students to hold urine, and 36% encourage holding by offering rewards to students who don't use bathroom passes or punishing those who do.<sup>4</sup> I frequently write letters on behalf of students whose bathroom access is restricted. During the coronavirus lockdowns, many of my patients' symptoms improved, largely because they could use the toilet without restriction.

Given all that, I have modified Dr. O'Regan's protocol in ways that have boosted my own success rates with his protocol. Some key modifications:

- 1.) I recommend daily enemas for at least 30 days, tapering only after the child remains dry for at least 7 consecutive days.** A significant number of children need daily enemas far longer than 30 days before tapering, especially those with both daytime and nighttime symptoms. I also recommend extending the second and third phases of **M.O.P.** to minimize the odds of relapse.
- 2.) I recommend a daily osmotic laxative in addition to enemas.** Whereas enemas do a powerful clean-out job, osmotic laxatives keep stool soft, helping children overcome their association of pooping with pain. Enemas work far better than laxatives alone, but the combination works best. However, for children with encopresis, osmotic laxatives often make matters worse to begin with, so I recommend adding these laxatives only after 2 weeks of daily enemas.
- 3.) After any 30-day period without progress, the protocol should be adjusted.** This can mean switching from phosphate enemas to liquid glycerin suppositories, shifting to large-volume enemas with glycerin and/or Castile soap, or adding overnight mineral-oil or olive-oil enemas and then moving the daily stimulant enema to the morning. The Anthology discusses these and other options.


3 Hodges S, Richards K, Gorbachinsky I, Krane LS. The association of age of toilet training and dysfunctional voiding. *Res Rep Urol*. 2014;6:127-130 <https://doi.org/10.2147/RRU.S66839>

4 Ko, L. N., Chuang, K., Champeau, A., Allen, I., & Copp, H. L. (2016, April). Lower Urinary Tract Dysfunction in Elementary School Children: Results of a Cross-Sectional Teacher Survey. *Journal of Urology*, 195(4), 1232-1238. doi:<https://doi.org/10.1016/j.juro.2015.09.091>

# Five Lessons from a Decade Prescribing M.O.P.

As I state in *The M.O.P. Book*, M.O.P. involves a lot of experimentation. I am always learning from the families I work with — and particularly from the thousands of parents who've cycled through the three private support groups I run on Facebook. These groups, which tend to attract the most challenging cases, serve as something of a laboratory, providing immediate feedback on various approaches. These families are the most desperate and the most willing to push the envelope, and I am grateful for all I learn from them.

Based on my experience with M.O.P., here are five lessons I would like to pass on to colleagues.



## 1.) Rectal diameter is critical

Dr. O'Regan noted that many parents had no idea their children were constipated, let alone severely so. This remains true. What's more, most of my patients have been referred by pediatricians who did not notice the patients were constipated or underestimated the severity of the constipation. I've had numerous patients with grapefruit-sized rectal masses, plainly visible via X-ray, that went unnoticed by the referring physician.

**This is because today's most common diagnostic methods — including the Bristol Stool scale, Rome criteria, patient history and exam — are of little value.** Many severely constipated children poop daily, but they don't fully evacuate. Palpating a child's abdomen is useless, too. Tiny, wiry kids can harbor massive amounts of stool that cannot be felt.

Constipation is often defined as "having fewer than 3 bowel movements a week," but this definition puts the emphasis on what is not happening (pooping) rather than what is happening: rectal stretching. Dr. O'Regan recognized this definition problem and made note of it in his studies.

*"When I started X-raying my patients and could see how much their rectums were distended, I realized enemas were the only thing that would make a difference."*

– James Sander, M.D.  
Director of Pediatric Urology, Doctors Hospital at Renaissance, Edinburg, Texas

The problem with relying on the frequency of bowel movements became obvious to me years ago, when I would refer enuresis cases to GI docs for evaluation. These physicians would refer the patients back to me, insisting these children were not constipated because they had normal marker studies. Yet when I scrutinized their X-rays, I saw rectums full of stool. I realized why: While some stool was passing through on a daily basis, it was oozing around the hard and growing mass that was aggravating bladder nerves.

Dr. O'Regan insisted on using anorectal manometry to detect constipation, knowing this method, though invasive, was the gold standard. Urology textbooks from 40 years ago echo his recommendations. But the



message got lost, and today's guidelines recommend a patient exam and pooping history. No wonder constipation routinely flies under the radar.

**It is also important to note that a severely constipated child may have a belly that looks and feels normal.** The rectum just stretches to accommodate the extra stool, and you just can't feel it.

**X-raying for constipation is not common medical practice, but for enuresis patients, it should be.** (There's no reason to X-ray for encopresis, since constipation is the only explanation.) We don't think twice about X-raying a child to diagnose a broken arm, an injury that will heal in 6 weeks. Yet many physicians won't X-ray to diagnose chronic constipation, a condition that can cause families years of suffering.

**Is it safe to X-ray for constipation? Yes.** The radiation dose of an abdominal X-ray is the same dose you get from simply living for 3 to 6 months. I'm not interested in putting my patients at risk for no good reason, and I am absolutely opposed to the overuse of X-rays in children. But the amount of good you can do for a child with bladder problems by accurately diagnosing constipation far outweighs the risks of a plain X-ray.

X-rays not only demonstrate to families why the child is wetting but also provide a baseline measurement for later comparison. Sometimes, when a child has been on **M.O.P.** for 30 days and shows no signs of progress, parents will say to me, "See, it wasn't constipation after all." An X-ray proves otherwise.

If you see an accumulation of stool in the rectum, no matter when the child last pooped, that is a problem. The rectum is designed to sense the arrival of stool and immediately empty, not to store it. It's not true that "there's always some poop in the rectum." **The most useful and precise way to assess constipation is to measure rectal diameter.** A measurement of greater than 3 cm at the rectum's widest point indicates constipation. Most of my enuresis patients have measurements 2 or 3 times normal.

## 2.) Enemas are far more effective than laxatives

Many of my patients have been prescribed Miralax — in many cases, huge amounts — but nonetheless ended up in my clinic because their enuresis did not resolve. I used to prescribe a lot of Miralax myself, until I recognized how inadequate it is. Often, it just makes matters worse, causing children to soil their pants.

For several years I relied on Dr. O'Regan's studies, as well as my own experience, to persuade families to give enemas a try. Eventually, colleagues and I conducted our own study, comparing the effectiveness of Miralax to enemas to treat enuresis.<sup>5</sup>

Our study tracked 60 patients, ages 4 to 11, with daytime enuresis. Forty patients followed standard therapies, including Miralax, timed voiding, and, in some cases, anticholinergics, and biofeedback physical therapy. Another 20 patients agreed to follow **M.O.P.**

**After 3 months, 30% of the patients treated with standard therapies had stopped wetting, compared to 85% in the M.O.P. group.**



5 Hodges, S. J., & Colaco, M. (2016). Daily Enema Regimen Is Superior to Traditional Therapies for Nonneurogenic Pediatric Overactive Bladder. *Global pediatric health*, 3, 2333794X16632941. <https://doi.org/10.1177/2333794X16632941>

A closer look at the data explains why enemas were so much more effective. At the start of the study, the average rectal diameter in both groups was greater than 6 cm. Three months later, the rectums of the 40 patients treated with standard therapy remained abnormally stretched — to 5 cm, on average. But rectal diameter in the enema group had been reduced to 2.15 cm, on average. Among the three **M.O.P.** subjects whose daytime enuresis persisted, rectal diameter remained abnormal. These patients were so stubbornly constipated that daily pediatric enemas were not sufficient to clean them out. (These are the kids who need the **M.O.P.+** regimen.)

In the generation since Dr. O'Regan's research was published, enemas have fallen out of favor, and I understand why. It's easy to hand a child a glass of water mixed with a tasteless, odorless powder. It's less easy for a parent to insert a tube up a child's bottom. But if you give families the options and the facts, most will choose "effective" over "easy."

**Osmotic laxatives do play a role in resolving enuresis, but only as a supplement to — not a substitute for — daily enemas. Stimulant laxatives can play an important adjunct role, too, as I explain in the Anthology.**

As for high-dose Miralax clean-outs, the results tend to be short-lived. I have had countless families go this route before ultimately conceding it wasn't working. They wish they'd tried enemas to begin with.

### 3.) Daily enemas do not cause dependence or electrolyte imbalance

Many of my patients have been told by pediatricians to avoid daily enemas "because they cause dependence and electrolyte imbalance." These warnings are not supported by research or by my experience, Dr. O'Regan's experience, or the experience of any other physician who uses **M.O.P.** or similar protocols. Let's explore these concerns.

**Dependence.** The notion that enemas cause dependence is commonplace but supported by no evidence I have come across. **I don't even know of a theoretical basis for the claim that enemas cause the bowel to stop working normally and that stool softeners are "safer."**

In a severely constipated child, the rectum is already not working normally. It has become so stretched that it has lost the ability to fully expel poop. **M.O.P.** allows the rectum to remain clear long enough to regain the sensation and tone necessary for full and complete emptying. Once the rectum has recovered, the child will no longer need enemas. One goal for a child on **M.O.P.** is to poop spontaneously once a day, in addition to pooping after each enema. If the child is only pooping after enemas, this is NOT a sign of dependence; it just means the child hasn't fully regained rectal tone and/or sensation.

**Electrolyte imbalance.** The concern about electrolytes pertains to over-the-counter pediatric enemas that contain phosphate. However, if you follow **M.O.P.** and FDA guidelines to limit enemas to one per day, this concern is unwarranted. Phosphate enemas are, of course, contraindicated for children with kidney disease, as I state in the **M.O.P.** safety guidelines.

Complications from enemas are so uncommon that a review of 39 studies conducted over 50 years found a total of only 15 cases of electrolyte imbalance in children ages 3 through 18.<sup>6</sup> Over 50 years. The vast majority of these cases involved children who had a chronic disease or were given more than one enema in a day.

6 MENDOZA, J., LEGIDO, J., RUBIO, S. and GISBERT, J.P. (2007), Systematic review: the adverse effects of sodium phosphate enema. *Alimentary Pharmacology & Therapeutics*, 26: 9-20. <https://onlinelibrary.wiley.com/doi/full/10.1111/j.1365-2036.2007.03354.x>

**In a decade of prescribing daily enemas, I have never had a patient develop an electrolyte imbalance.** If this were a legitimate concern, I would have found out by now.

Nonetheless, anyone concerned about electrolyte imbalance can easily implement **M.O.P.** with alternative enemas — such as liquid glycerin suppositories or large-volume enemas with glycerin — that pose no risk whatsoever of electrolyte imbalance. I discuss various options in *The M.O.P. Book: Anthology Edition*.

#### 4.) Enemas are not “abusive” or “traumatic”

Unlike physicians in Dr. O'Regan's day, the medical community today has deemed enemas overly aggressive, if not “abusive,” a term many of my patients have heard from other physicians. But this notion runs contrary to the experience of the families I work with. No studies have considered whether a regimen such as **M.O.P.** will “emotionally scar” a child, as one doctor put it, but I cannot find any basis for this concern.

Certainly many parents and children are apprehensive, if not downright fearful, when they start enemas. Mostly they fear enemas will hurt, and sometimes they do hurt. This is usually because the child isn't relaxed or positioned correctly or because there's not enough lubrication on the tip. I address these issues in a blog post titled “[13 Ways to Ease Your Child's Fear of Enemas.](#)”

For most families, enemas quickly become routine, even something to look forward to. Yes, that's true! As one mom in our support group posted: “My 4.5-year-old loves his enemas! He was constipated and had stomach aches for over 2 years with nothing else helping. Now he's going on his own, dry and clean all day and no tummy aches.”

Another mom posted that it was oral laxatives, including Miralax, that caused her daughter distress: “Cleanouts gave her messy accidents, humiliation, and painful rashes. My daughter loves her enemas!”


I hear this all the time. Yet many physicians find it impossible to believe children are fine with enemas. One mom told me: “When my child told the doctor she liked enemas because she felt better, her statement was immediately dismissed with a ‘No, you don't. No one likes them.’ ”

Another mom wrote: “Our daughter's urologist still makes a stinky face and says it's a really aggressive treatment choice.”

Yes, **M.O.P.** is aggressive — in the best possible way. **Chronic constipation in children is a notoriously stubborn problem, and in my experience, aggressive treatment is the only kind of treatment worth doing.**

*“My daughter likes how she feels after her enemas. Does she want them to end one day? Yes! But she is perfectly content because she has seen how getting them helps. And it helped her to be comfortable pooping at school because she can feel the urge now.”*

— Mother of **M.O.P.** patient



## 5.) Alternatives to Miralax can work just as well

In my experience, Miralax is the easiest and most effective osmotic laxative to give children as part of **M.O.P.** However, a growing number of parents are expressing concern about Miralax safety, and some members of our support group have reported behavioral or psychiatric disturbances in their children after starting PEG 3350. I do not push Miralax on these or any other parents. Lactulose works equally well for many children, and some parents report their children like it better. Many parents use magnesium citrate or magnesium hydroxide (milk of magnesia) with good results. With so many effective alternative laxatives on the market, there is no reason to steer parents toward Miralax if they have concerns.

# The M.O.P. Progression

The Modified O'Regan Protocol is a trial-and-error process. If your child goes 30 days without progress, make a change!

## – Store-bought enema + osmotic laxative

**M.O.P.**

### Enema options:

- phosphate enema (pediatric or adult size)
- liquid glycerin suppository (pediatric or adult size)



### Laxative options:

- PEG 3350 (Miralax)
- magnesium citrate
- lactulose
- magnesium hydroxide

OPTIONAL: Daily or periodic stimulant laxative for kids who haven't achieved daily spontaneous poop.

## – Large-volume enema + osmotic laxative

**M.O.P.+**

### Enema solution progression:


First, try saline + glycerin. Then...



...add Castile soap. Then...



...add phosphate enema contents.



OPTIONAL: Daily or periodic stimulant laxative for kids who haven't achieved daily spontaneous poop.

## – Oil-retention enema (evening) + large-volume or store-bought enema (morning) + osmotic laxative

**DOUBLE M.O.P.**

Oil-retention enemas can be done daily, weekly, or periodically.

Oil options: mineral oil



olive oil



## – Smaller enema volume + stimulant laxative

**M.O.P.x**

### Enema options:

- reduce homemade saline solution while maintaining stimulant dose
- return to store-bought enema or LGS



+




NOTE: Use M.O.P.x when daytime accidents have stopped on M.O.P.+ but bedwetting persists

# Resources for Parents





Our website, [BedwettingAndAccidents.com](http://BedwettingAndAccidents.com), offers numerous books, guides, videos, and infographics you can use to educate patients. I am happy to share, at no cost, the PDF version of any of our books. Here's an overview of what we offer.

Here's an overview of what we offer. I have attached some of our infographics.


## Books for Parents



## Books for Kids



## Infographics



# 12 Signs

## Your Child

# is Constipated

Holy Cow!

- 1** **XXL poops.** We're talking "Holy cow!" poops – larger than 3/4" x 6."

- 2** **Firm poops.** Logs or pellets = bad; thin snakes or mushy blobs = good.



- 3** **Poop accidents.** When the rectum is overstuffed, poop just falls out.



- 4** **Bedwetting and pee accidents.** A big 'ol poop mass aggravates the bladder.

AGAIN?  
But you JUST Peed!

- 5** **Recurrent UTIs.** Bacteria from overflowing poop crawl up to the bladder.


- 6** **Extremely frequent and/or urgent peeing.** You think, "AGAIN? But you JUST peed!"

- 7** **Infrequent pooping.** But daily pooping doesn't rule out constipation.

- 8** **Pooping more than 2x/day.** A stretched-out rectum lacks the tone to evacuate fully.

- 9** **Belly pain.** Constipation is the #1 source of tummy ache in kids.


- 10** **Skid marks or itchy anus.** Clogged kids can't fully empty  
→ bottom is hard to wipe  
→ poop stains.



- 11** **Super-loose poop.** Some poop can ooze around the large, hard rectal clog.




- 12** **Continued trouble toilet training.** Your child may fear pooping or hide to poop in diapers.



BedwettingAndAccidents.com

Regan Press

# How's Your Poop?




Smooth  
mushy  
snakes



mushy  
blobs




soft-  
serve  
ice cream



Healthy  
POOP

Oh  
No!




Clogged  
Colon

© CRISTINA ACOSTA

rabbit  
pellets



big ol'  
logs



thick &  
bumpy  
sausage





# Bedwetting: 4 Truths

Every Parent  
Must Know

**TRUTH  
#1**

Some children don't  
outgrow bedwetting.

**70%:** Odds a bedwetting 9-year-old, left untreated, will wet the bed at 19 <sup>[1]</sup>

**840,000:** Number of U.S. tweens/teens who wet the bed <sup>[2]</sup>

**32:** Percent of bedwetting teens/tweens who also have daytime accidents <sup>[3]</sup>

Bedwetting is less likely to resolve spontaneously if a child:

- wets the bed nightly
- also has daytime pee or poop accidents



Bedwetting is  
**NOT** caused by:

~~Deep sleep~~  
~~Stress or laziness~~  
~~Underdeveloped bladder~~  
~~Urine overproduction~~

**TRUTH  
#2** Bedwetting is caused by  
chronic constipation.

Poop piles up and stretches the rectum, which presses against and aggravates the bladder. X-rays prove it. When constipation resolves and the rectum shrinks back to size, research shows, bedwetting stops. <sup>[4]</sup>

**TRUTH  
#3** Bedwetting should be  
treated around age 4.

Waiting until age 7 does not serve the child. The longer the rectum remains clogged and stretched, the more difficult the fix.

**TRUTH  
#4** Enemas work far better  
than alarms or medication.

Medication has a dismal success rate and, like an alarm, does nothing to resolve the root cause: constipation. A regimen of enemas + laxatives will resolve bedwetting for good.

Take action today! Learn more at  
**BedwettingAndAccidents.com**

© Steve Hodges and Suzanne Schlosberg 2017  
Design by DyanRothDesign.com

[1] <http://onlinelibrary.wiley.com/doi/10.1111/j.1464-410X.2006.06074.x/full>

[2] <http://www.actforyouth.net/adolescence/demographics/>

[3] <http://onlinelibrary.wiley.com/doi/10.1111/j.1464-410X.2006.06074.x/full>

[4] <https://www.bedwettingandaccidents.com/research>

# 4 MiraLAX Facts You Didn't Know

The real poop on giving children PEG 3350

## FACT #1

### Doctors rely too heavily on MiraLAX.

The problem with PEG 3350: Soft poop often oozes around the clog, solving nothing. Chronically constipated kids also need enemas or glycerin suppositories (solid or liquid).

WHAT MIRALAX DOES:

Draws water into the colon, softening stool so pooping doesn't hurt.

WHAT MIRALAX DOESN'T DO:

Dislodge the hard, rectum-clogging stool mass that causes bedwetting and accidents.

## FACT #2

### Enemas resolve wetting far better than MiraLAX.

In a study of 60 children\* who wet their pants daily, after 3 months:



of children treated with PEG 3350 stopped wetting



of children treated with enemas stopped wetting

Why? Because enemas let the rectal diameter shrink back to normal size: < 3 cm.

After 3 months, rectal diameter of:

- PEG 3350 group: 5 cm
- Enema group: 2.15 cm



## FACT #4

### MiraLAX is **not** a lifetime solution to constipation!

Once constipation resolves, children should stay on osmotic laxatives for 3 to 6 months and must:

- Eat "real" rather than highly processed food
- Drink plenty of water
- Stay active
- Poop with feet on a tall stool


## FACT #3

### If you worry about PEG 3350 safety, you have options.

Effective osmotic laxatives with unquestioned safety include:

- Magnesium citrate (powder or capsules)
- Lactulose (prescription liquid)
- Magnesium hydroxide (chewable tablets)

FDA-funded researchers are investigating whether PEG 3350 triggers psychiatric symptoms, as some parents have reported. So far, 100+ studies have found PEG 3350 safe for children.



A hilarious book about healthy pooping.

[BedwettingAndAccidents.com](http://BedwettingAndAccidents.com)

# 9

**Kid-Tested**



# WAYS TO MAKE ENEMAS LESS SCARY

**Compare the enema tip to a typical poop.**



Constipated kids' jumbo stools are *far* wider than an enema tip!

**Let your child take control.**

Young ones can open the package and take the cap off; older kids can give themselves the enema.

**Help your child relax.**

Encourage deep breathing, like blowing out birthday candles.



**Offer a small reward.**

Screen time, a Starburst, a dollar – whatever works!

**Do a demo.**

Squirt a sugar cube with a syringe of water until it starts getting mushy. Kids love the visual.



**Give yourself an enema.**

Yes, you! What better way to show empathy and offer a scouting report?

**Read "Bedwetting & Accidents Aren't Your Fault."**

Your child will know just what to expect.



**Add lubrication.**

K-Y Jelly or Vaseline on the tip and/or your child's bottom will ease the way.

**Try different brands or ingredients.**

To some kids, the shape of the tip matters. To others, liquid glycerin suppositories are gentler.

# Full Text of Key Studies

## Link between constipation and enuresis, encopresis, UTIs (the O'Regan studies):

- [Relevance of constipation to enuresis, urinary tract infection and reflux. A review.](#)  
Yazbeck S, Schick E, O'Regan S., Eur Urol, 1987
- [Constipation, bladder instability, urinary tract infection syndrome.](#)  
O'Regan S, Yazbeck S, Schick E., Clin Nephrol, 1985
- [Constipation a commonly unrecognized cause of enuresis.](#)  
O'Regan S, Yazbeck S, Hamberger B, Schick E., Am J Dis Child, 1986
- [Constipation and the Urinary System](#)  
Sean O'Regan, Salem Yazbeck and Eric Schick, "Pediatric Urology," 1997; Chapter 16, pp. 197-199

## Effectiveness of enemas v. Miralax for treating enuresis (comparison of rectal diameter):

- [Daily Enema Regimen Is Superior to Traditional Therapies for Nonneurogenic Pediatric Overactive Bladder](#)  
Hodges SJ, Colaco M. Global Pediatric Health, 2016, 3: 1-4

## Safety of enemas in children:

- [Systematic review: the adverse effects of sodium phosphate enema.](#)  
Alimentary Pharmacology and Therapeutics. Mendoza, J., Legido, J. 22 April 2007

## Evidence that bedwetting will not spontaneously resolve in some children:

- [Differences in characteristics of nocturnal enuresis between children and adolescents: a critical appraisal from a large epidemiological study.](#)  
Yeung, C., Sreedhar, B., et al. BJUI International. 6 April 2006

## Relevance of Constipation to Enuresis, Urinary Tract Infection and Reflux

### A Review

*Salam Yazbeck, Erik Schick, Sean O'Regan*<sup>1</sup>

Department of Pediatric Nephrology and Surgery, Centre de Recherche Pédiatrique, Hôpital Ste-Justine, University of Montreal, Que., Canada

**Key Words.** Constipation · Urinary tract infection · Enuresis

**Abstract.** Little attention has been afforded the relevance of constipation to urinary symptoms. Problems of definition and measurement have contributed to this problem. A review of the literature suggests that rectal dilatation may influence the function of the urinary tract leading to urinary tract infection and enuresis. Our studies indicate that constipation may, by causing uninhibited bladder contractions, cause urinary tract infection, enuresis and vesicoureteral reflux.

The world of medicine has undergone revolutionary changes, with quantum leaps in information on various organ systems being realized in the last 3 decades. As a result, not only has specialization evolved concentrating on single organ systems, but the explosion of knowledge has been such that subspecialization based on elements within organ systems is commonplace. Attention by specialists is usually focused on adjacent organ systems only when infiltrative lesions are involved, distant derangements occur due to hormonal effects or toxin accumulation or when multi system organ dysfunctions occur due to systemic disease. Consequently in the field of nephro-urology relatively scant attention has been afforded to observations relating derangements in bowel function which might influence the function of the urinary tract system.

During the past 5 years we have focused our attention in our nephro-urology clinics to determining the relevance of constipation to pediatric problems associated with enuresis, urinary tract infection and vesicoureteral reflux. We review evidence substantiating a role for constipation in causing urinary symptoms.

Prior to puberty the pediatric abdominal and pelvic cavities are one. As a result the close proximity of the rectum to the posterior wall of the bladder is such that any gross distention of the rectum could therefore result

in compression of the bladder, bladder neck obstruction or distension of the urethra leading to abnormalities of urinary tract function [1]. Shoppner [2] in 1968 demonstrated that the presence of constipation has the potential for inducing gross distortion of the bladder and urethra in children. Though this report concentrated on radiologic evidence of distortion resulting from constipation, of note was the fact that of the 39 children studied, 8 had reflux, 21 also had enuresis and 2 had recurrent urinary tract infection. The relevance of these abnormalities to the constipation was not discussed. Neumann et al. [3] in 1973 noted the association of constipation and recurrent urinary tract infection in children. On occasion they also noted the presence of reflux in these constipated children and they indicated that the aggressive treatment of constipation resulted with its resolution of associated urinary tract infection. In addition these workers made the point that constipation was not sought commonly as a complaint because: (a) the doctors did not inquire; (b) bowel preparation prior to radiologic studies resulted in evacuation, eliminating radiologic signs of constipation, and (c) mothers did not volunteer the information because of either lack of

<sup>1</sup> Supported by a grant from the Hospital for Sick Children Foundation (Toronto) and NATO.

knowledge of the presence of constipation or because it was not a dominant presenting complaint.

Progress in this field has been hampered by the lack of a standard definition of constipation. Also a method of measurement of rectal abnormality which would correlate with the presence of constipation and which was recordable and therefore free of observer bias was not generally available.

Progress in the past decade has enabled the development of methods and definitions which go some way to achieving these ends. We [4] therefore define functional constipation as being present if any of the following are present: (1) more than a 72-hour interval between bowel movements; (2) the presence of an overflow fecal incontinence (soiling or encopresis); (3) the passage of small hard scibalous stools with intermittent passage of large stools; (4) poor emptying and dilatation of rectal ampulla after defecation as determined by rectal examination, and (5) grossly decreased level of perception and increased tolerance of balloon insufflation in the presence of normal anorectal relaxation during rectal manometry, combined with any element of the above.

We performed rectal manometric studies using an air-filled balloon system. We noted the smallest inflation volume perceived by the patients, the volume at which relaxation of the internal sphincter occurred and the maximum inflation volume that was tolerated by the patients without pain or discomfort [5]. The diagnosis of constipation is a diagnosis to be made by a physician with appropriate questioning (quantity and quality of stools, frequency of defecation, etc.) and physical examination. A convenient way of avoiding being misled is to obtain a written record of stool frequency and quality at follow-up visits.

Similarly apart from radiologic studies, acceptable measurements and recording of bladder and urethral function under normal and abnormal conditions were generally not available. However, the advent of techniques to assess urodynamics has led to the identification of uninhibited bladder contractions as a common finding in children with urinary tract infection. The presence of uninhibited bladder contractions were determined by urodynamic studies using a DISA 1200 system [4]. Bladder instability was considered present if at least one of these elements were identified: (1) the presence of uninhibited contractions of the detrusor during the filling phase of the bladder with an amplitude equal or greater than 15 cm of water and (2) the occurrence of detrusor contraction at the end of or after urinary flow.

### Constipation and Urinary Tract Infection

Using these criteria for constipation and uninhibited bladder contractions we studied 47 children with recurrent urinary tract infections [6]. In all these patients follow-up urodynamic studies indicated uninhibited bladder contractions. They were also constipated. Enuresis was present in 32 and encopresis in 21. Aggressive treatment of the constipation resulted in cessation of infection in 44 of the 47 children, enuresis in 22 of 32 patients and encopresis in 20 of 21 patients and an improvement of bladder function as indicated by cessation of uninhibited contractions in 12 who underwent repeat studies. These results were achieved by appropriate dietary advice (increase fiber content) and by aggressive treatment of constipation utilizing a phosphate soda enema daily for 1 month, every other day for a further month and twice weekly or more often if required for a third subsequent month. This was in order to allow for return of normal rectal tone by a dilated hypotonic rectum. Of interest was that many of the patients studied had no prior history, or a history of constipation was denied. Normal bowel habits were perceived by patients and parents even in the presence of proven rectal reservoirs of feces by rectal examination and rectal manometry and on occasion even with encopresis. White and Taylor [7] have also noted the frequent presence of constipation with urinary tract infection as have Smellie et al. [8]. A retrospective analysis of the incidence of urinary tract infection in patients with Hirschsprung's disease, a constipating entity, disclosed a much increased frequency of infection prior to and after surgery [9]. An increased incidence of bacteruria occurs in rats with fecal retention [10].

### Constipation and Enuresis

Therapy for enuresis presents a major problem for pediatric nephrologists and urologists. Psychologic stress inflicted on the patients by discomfort of bed wetting and by parental reaction is considerable. As a result numerous therapeutic modalities have been used including alarm devices and systemically acting drugs, none of which is completely satisfactory [11]. The exact etiology of enuresis is unknown. However, we did note a very high incidence of enuresis in girls with urinary tract infection and constipation [6]. The enuresis resolved upon treatment of constipation. We therefore studied 22 patients with enuresis [4]. We noted that in over 40% of these cases encopresis or soiling was also present but was considered a

minor symptom. By history, physical examination including rectal examination and rectal manometric studies, constipation was an extremely common though often unrecognized accompaniment of enuresis. These patients had uninhibited bladder contractions similar to that previously observed by Berger et al. [12] who also noted a decreased bladder capacity in relation to age in many of these patients, a phenomenon similarly noted by us. Aggressive treatment of constipation resulted in rapid resolution of enuresis without any form of pharmacologic drug therapy. Our studies strongly implicate unrecognized rectal distension as an etiologic factor in enuresis.

As previously stated, Shopfner [2] in discussing urinary tract pathology associated with constipation also noted that 54% of these patients were enuretic. Similarly Baumann and Hinman [13] described the treatment of incontinent boys with nonobstructive disease of the urinary tract. These children had encopresis and enuresis. Hypnotherapy was emphasized for the therapy of the enuresis. However, they did note that aggressive treatment of constipation including, if necessary, digital evacuation of the rectum was required prior to cessation of enuresis. This fact suggests to us that the constipation was the primary element in causing enuresis. We have also noted uninhibited bladder contractions in children who were constipated but did not have urinary symptoms [4]. Uninhibited bladder contractions in the absence of urinary incontinence can occur in situations in which urethral constriction can overcome the increased intravesical pressures induced by uninhibited contractions [14]. The presence of uninhibited contractions in otherwise asymptomatic but constipated children [4] suggests a cause and effect relationship. Drug therapy directed towards the suppression of uninhibited bladder contractions in enuretic [15] or refluxing patients [16] may be successful. However, aggressive treatment of constipation, a therapy directed against a common etiologic factor causing bladder abnormalities, may be more beneficial.

### **Constipation with Vesicoureteral Reflux**

Though little attention has been paid to the observation, constipation was noted to cause dilatation of the urinary system [16]. Dilatation of the urinary tract, especially the bladder, but also the ureters was noted by numerous authors to occur in association with Hirschsprung's disease [17-19]. In addition successful treatment of Hirschsprung's disease may result in resolution of the vesicoureteral reflux [20]. Kottmeier and Clatt-

worthy [21] noted a similar incidence of vesicoureteral reflux in children with severe functional constipation and those with Hirschsprung's disease. We noted constipation to be present in patients with primary vesicoureteral reflux [22]. Though numbers did not allow for a controlled study, we did note rapid resolution of reflux with aggressive treatment of constipation in the absence of antibiotic therapy, or anticholinergic therapy for treatment of associated inhibited bladder contractions. Similarly White [23] noted that resolution of infection with reflux is more easily attained when accompanying constipation is aggressively treated. These reports suggest that constipation is a nonfortuitous phenomenon occurring in patients with vesicoureteral reflux and may be of major etiologic importance. Consequently this phenomenon may be a major factor of contention in the interpretation of the medical-surgical prospective study of the International Study on Vesicoureteral Reflux since, in the medical protocol, avoidance of constipation is emphasized whereas this factor is not considered in the surgical approach to therapy [24].

Hinman [25] made note of the fact that the constipated child may have an evolution of encopresis leading to enuresis, to urinary tract infection and eventually to vesicoureteral reflux. Though the symptom complex was attributed by him to behavioral characteristics, perhaps constipation may be the initiating factor as resolution of reflux and enuresis followed therapy including treatment of encopresis [26]. The precipitating cause of constipation may be due to short episodes of psychologic stress (e.g., to early aggressive toilet training, to anal fissure, etc.), causes that may be long resolved but which, however, may result in persisting constipation.

Studies by Bailey et al. [27] noted a 55% incidence of abnormal anal sphincter electromyograms in children with enuresis and urinary tract infection. These observations substantiate the possibility that abnormalities of the rectum may cause enuresis, urinary tract infection and reflux in children. The observation that children with functional constipation may have uninhibited contractions of the bladder in the absence of urinary symptoms strongly supports this possibility. Because the rectal sphincter and the urethral sphincter, together with the perineal musculature is considered as a single physiologic unit, with voluntary rectal sphincter contraction, consequent concomitant urethral sphincter contraction occurs. This urethral sphincter contraction, in turn, may be responsible for a dyssynergistic voiding pattern with secondary bladder instability, enuresis, urinary tract infection, and/or vesicoureteral reflux possibly ensuing. Thus a

spectrum of disease may exist of constipation with uninhibited bladder contractions in the absence of urinary symptoms or with enuresis, urinary tract infection and vesicoureteral reflux. Because of the high spontaneous resolution rate of vesicoureteral reflux, a long-term controlled trial of constipation treatment would be required to substantiate the thesis of a cause and effect relationship.

Since the intestinal transit time in children with functional constipation is normal, oral therapy alone directed to maintenance of an empty rectum is inappropriate [28]. Though we have achieved satisfactory success using phosphate soda enema therapy, factors of cost as well as patient discomfort and inadequate volume to achieve consistent and complete rectal evacuation have arisen. A more appropriate and satisfactory approach is the use of saline enemas (5 ml of salt in 1 liter of warm water). Using a bag held 2 feet above the patient an adequate volume will be infused under gravity. The incidence of cramps is less with this method [Murray, R., personal commun.].

The morbidity associated with enuresis, urinary tract infection and reflux is so high and constipation a condition easily amenable to resolution with aggressive therapy that extreme care should be exercised in determining whether constipation is present in children presenting with these complaints, so that therapy for constipation may be instituted and thus aid in the resolution of the urinary symptoms.

## References

- O'Regan, S.; Yazbeck, S.: Constipation: a cause of enuresis, urinary tract infection and vesico-ureteral reflux in children. *Med. Hypotheses* 17: 409-413 (1985).
- Shopfner, C.E.: Urinary tract pathology associated with constipation. *Radiology* 90: 865-877 (1968).
- Neumann, P.Z.; de Domenico, I.J.; Nogrady, M.B.: Constipation and urinary tract infection. *Pediatrics, Springfield* 52: 241-145 (1973).
- O'Regan, S.; Yazbeck, S.; Hamberger, B.; Schick, E.: Constipation a commonly unrecognized cause of enuresis. *Am. J. Dis. Child.* 140: 260-261 (1986).
- Meunier, P.; Mollard, P.; Marechal, J.M.: Physiopathology of mega rectum: the association of megarectum with encopresis. *Gut* 17: 224-227 (1976).
- O'Regan, S.; Yazbec, S.; Schick, E.: Constipation, bladder instability, urinary tract infection syndrome. *Clin. Nephrol.* 23: 152-154 (1985).
- White, R.H.R.; Taylor, C.M.: The non-operative management of primary vesicoureteric reflux; in Johnson, Management of vesicoureteric reflux, pp. 117-136 (Williams & Wilkins, Baltimore 1984).
- Smellie, J.M.; Katz, G.; Grüneberg, R.N.: Controlled trial of prophylactic treatment in childhood urinary tract infection. *Lancet* ii: 175-178 (1978).
- Yazbeck, S.; O'Regan, S.: Hirschsprung's disease and urinary tract infection: unrecognized association. *Nephron* 43: 211-213 (1986).
- Breda, G.; Bianchi, G.P.; Bonimi, U.; Piacentini, I.; Farello, G.: Faecal stasis and bacteriuria: experimental research in rats. *Urol. Res.* 2: 155-157 (1975).
- Kass, E.J.; Diokno, A.C.; Montealergre, A.: Enuresis: principles of management and result of treatment. *J. Urol.* 121: 794-796 (1979).
- Berger, R.M.; Maizels, M.; Moran, G.C.; Conway, J.J.; Firlit, C.F.: Bladder capacity (ounces) equals age (years) plus 2 predicts normal bladder capacity and aids in diagnosis of abnormal voiding patterns. *J. Urol.* 129: 347-349 (1983).
- Baumann, F.W.; Hinman, F.: Treatment of incontinent boys with nonobstructive disease. *J. Urol.* 111: 114-116 (1974).
- Koff, S.A.: Disordered vesico-urethral function in the pathogenesis of urinary infection and vesico-ureteric reflux; in Johnson, Management of vesicoureteric reflux, pp. 67-81 (Williams & Wilkins, Baltimore 1984).
- Smey, P.; Firlit, C.F.; King, L.R.: Voiding pattern abnormalities in normal children. Results of pharmacologic manipulation. *J. Urol.* 120: 574-577 (1978).
- Koff, S.A.; Murtagh, D.S.: The uninhibited bladder in children: effect of treatment of recurrence of urinary tract infection and on vesico-ureteral reflux resolution. *J. Urol.* 130: 1138-1141 (1983).
- Savage, J.P.: The deleterious effect of constipation upon the reimplanted ureter. *J. Urol.* 109: 501-503 (1973).
- Swenson, O.; Fisher, J.M.: The relationship of megacolon and megaloureter. *New Engl. J. Med.* 253: 1147-1150 (1955).
- Ehrenpreis, T.: Hirschsprung's disease, pp. 60-61 (Year Book Medical, Chicago 1970).
- Sieber, W.K.; Soave, F.: Hirschsprung's disease. *Curr. Probl. Surg.* 15: 14-15 (1978).
- Kottmeier, P.K.; Clattworthy, D.W.: Aganglionic and functional megacolon in children - a diagnostic dilemma. *Pediatrics, Springfield* 36: 572-582 (1965).
- O'Regan, S.; Schick, E.; Hamberger, B.; Yazbeck, S.: Constipation associated with vesico-ureteral reflux. *Urology* 28: 394-396 (1986).
- White, R.: Reflux nephropathy. Update: 1983. Discussion. *Contr. Nephrol.*, p. 249 (Karger, Basel 1984).
- Medical versus surgical treatment of primary vesicoureteral reflux: A prospective international reflux study in children. *J. Urol.* 125: 277-283 (1981).
- Hinman, F.: Urinary tract damage in children who wet. *Pediatrics, Springfield* 54: 143-150 (1974).
- Hinman, F.; Bauman, F.W.: Vesical and ureteral damage from voiding dysfunction in boys with neurologic or obstructive disease. *J. Urol.* 109: 727-732 (1973).
- Bailey, J.A.; Powers, J.J.; Wayeonis, G.W.: A clinical evaluation of electromyography of the anal sphincter. *Archs phys. Med. Rehabil.* 51: 403-408 (1970).
- Corazziari, E.; Cucchiara, S.; Staiano, A.; Romaniello, G.; Tamburrin, O.; Torsoli, A.; Auricchio, S.: Gastrointestinal transit time, frequency of defecation, and anorectal manometry in healthy and constipated children. *J. Pediat.* 106: 379-882 (1985).

Sean O'Regan, MD, Department of Pediatric Nephrology  
Centre de Recherche pédiatrique, Hôpital Ste-Justine  
3175 Côte Ste-Catherine, Montreal, Que. H3T 1CS (Canada)



# Constipation, bladder instability, urinary tract infection syndrome

S. O'REGAN, S. YAZBECK and E. SCHICK

Department of Pediatrics, Université de Montréal, Centre de Recherche Pédiatrique, 3175, Chemin Côte Sainte-Catherine, Montréal, Québec, Canada H3T 1C5

**Abstract.** Forty-seven children with recurrent urinary tract infection were noted to have large fecal reservoirs by rectal examination and rectal manometry. Constipation was accompanied in the majority by enuresis and/or encopresis. Urodynamic studies indicated uninhibited bladder contractions. Aggressive treatment of the constipation resulted in cessation of infection in 44 patients, enuresis in 22 of 32 patients and encopresis in 20 of 21 patients and an improvement in bladder function with cessation of all other forms of treatment.

**Key words:** urinary tract infection – constipation – bladder contractions

## Introduction

Recent studies have indicated that some children may have urodynamic abnormalities associated with recurrent urinary tract infections in the absence of radiologic abnormalities of the urinary tract [Allen and Bright 1978, Firlit et al. 1978, Smey et al. 1978]. It has been indicated in one study that abnormal bowel habits as determined by parental response to questioning may be observed in some patients with urinary tract infection [Neumann et al. 1973]. Distortion of the urinary tract system associated with constipation, though in the absence of urinary tract infection has also been reported [Shopfner 1968]. Also anal electromyography has been reported as abnormal in 57% of children with urinary tract infection [Bailey et al. 1970]. We describe the association of constipation, commonly unrecognized, with uninhibited bladder contractions as determined by urodynamic studies in children with symptomatic urinary tract infection without radiologic evidence of anatomic abnormality.

## Patients and methods

The identification of several children investigated for recurrent urinary tract infection who had uninhibited bladder contractions and functional constipation, all of whom had large fecal reservoirs as determined by rectal examination and rectal manometry,

led to the investigation of 47 patients who had no evidence of radiologic abnormality but had a similar pattern of abnormal bladder contractility and recurrent urinary tract infection. All were girls, with a mean age of  $8.2 \pm 2.53$  years (1 SD) and a mean duration of symptoms of  $3.7 \pm 2.28$  years (1 SD). Mean age of onset of first urinary tract infection was  $4.6 \pm 2.26$  years. Close questioning of the parents of several children did disclose elements suggesting the presence of chronic constipation. However, in 21 cases, constipation was denied as a symptom. All patients had been referred for assessment because of culture proven recurrent urinary tract infection with (32 patients) or without (15 patients) enuresis. Twenty one patients had mild encopresis. Radiologic investigation in all had been negative. Three patients had an anterior anus as determined by a decreased distance between the anal orifice and the vestibule. Because of the identification of the typical urodynamic tracings of an unstable bladder, later patients underwent directly rectal manometry in the absence of a history of constipation or encopresis. All patients had normal renal function as determined by normal plasma creatinine levels.

*Urodynamic studies were performed using a DISA 2100 Urosystem*

The patient was placed in a lithotomy position. After disinfection and draping, a 7-F single microtip transducer catheter was passed into the bladder per urethram and the bladder was filled. A balloon filled with water was placed in the rectum to register intra-abdominal pressure variations. Circular surface electrodes were placed on each side of the anus, and were used for perineal electromyography.

Received September 2, 1984, in revised form October 11, 1984.

Reprint requests to Dr. Sean O'Regan.

This paper was supported by a grant from the Hospital for Sick Children Foundation.

The child was seated on a flowmeter chair and data were recorded on a six channel recorder. The recorder noted simultaneously the total bladder pressure, the intra-abdominal (rectal) pressure and the intrinsic bladder pressure, obtained by the subtraction of the rectal pressure from the total bladder pressure. The bladder was filled with NaCl 0.9% at room temperature and 2 to 3 complete voiding cycles were registered on each child.

Bladder instability was considered present if at least one of these elements was identified:

1. The presence of uninhibited contractions of the detrusor during the filling phase of the bladder with an amplitude equal or greater than 15 cm H<sub>2</sub>O.
2. The occurrence of detrusor contraction at the end of or after urinary flow.

#### Rectal manometry performed as follows

Rectal manometric studies were done using a Beckman 710 recorder with a Shuster air filled balloon system [Menuier et al. 1967]. We noted the smallest inflation volume perceived by the patient, the volume at which relaxation of the internal sphincter occurred and the maximal inflation volume that was tolerated by the patient without pain or discomfort.

Functional constipation was deemed present when the following elements were noted:

1. Decreased perception and increased tolerance in response to large volume stimulation by the rectal balloon.
2. The presence of a normal rectal-anal reflux.

Twelve patients had been treated with oxybutyryn chloride (Ditropan) to inhibit abnormal bladder contractions without satisfactory response. All these patients with functional constipation were treated with a phosphate-soda enema (Fleets) once a day for one month and once every other day for the 2nd month to maintain the dilated rectum empty and prevent re-accumulation while awaiting the return to a normal rectal tone. All other forms of treatment were stopped [oxybutyryn chloride (Ditropan, antibiotics)] when the enema treatment was initiated.

#### Results

A consistent pattern of uninhibited bladder contractions was observed in all patients studied, consisting of detrusor contraction during the filling phase of the bladder with an amplitude equal or greater than 15 cm H<sub>2</sub>O or the presence of detrusor contraction at the end of or after urinary flow [Allen and Bright 1978, Firlit et al. 1978].

Though sensation under normal circumstances can be determined when the balloon of the rectal




Fig. 1 Bladder capacity in 36 children with urinary tract infection and constipation. Upper and lower lines indicate 95% confidence limits [Berger et al. 1983].

manometer is inflated with 5 ml to 10 ml of air the majority of these children had poor perception of rectal distension until at least 40 ml had been instilled into the balloon. All patients could tolerate balloon distension of 80 to 110 ml (maximal balloon insufflation volume; 110 ml, 6.7 cm diameter) without experiencing any discomfort. A majority of patients had a decrease in urinary bladder capacity. Three patients were noted to have an anterior anus.

Dramatic improvement was evident in all patients who followed the enema regimen. At follow-up clinic visits during which cultures were done only 3 patients had further episodes of urinary tract infection after initiation of treatment. At follow-up, constipation persisted in 2 patients. Enuresis ceased in 22 patients and improved (1 night/wk to month) in 7. Encopresis resolved in 20 patients. Two of 3 patients who did not respond completely were noted to have refused to follow the enema regimen appropriately. At follow-up of 17 to 9 months (mean  $12 \pm 2$ ), elimination of constipation resulted in dramatic symptomatic and psychologic improvement in 44 patients. In 12 patients in whom follow-up urodynamic studies were done, normal bladder function as determined by urodynamic studies has been attained. Control rectal manometry was performed in some patients and showed perception to lower volumes of balloon insufflation.

#### Discussion

Functional constipation has been associated with radiologic distortion of the urinary system in the presence and absence of urinary tract infection

[Neumann et al. 1973, Shopfner 1968]. We describe the association of rectal reservoirs of feces and uninhibited bladder contractions with enuresis, encopresis and recurrent urinary tract infection. In many of the patients studied there was no prior history or a history of constipation was denied. Normal bowel habits were perceived by patients and parents even in the presence of proven rectal reservoirs of feces by rectal examination and rectal manometry and on occasion even with encopresis. Close questioning however, indicated that patients had been toilet-trained very early in childhood and either had large infrequent stools or incomplete evacuation as determined by rectal examination after defecation. Elimination of constipation resulted in dramatic symptomatic improvement with cessation of urinary tract infection and enuresis even in the presence of some continuing abnormal bladder contractility.

Uninhibited bladder contraction is typical of that seen in infancy before full maturation of cortico-spinal control of bladder activity is achieved [Smey et al. 1978]. It is possible that the presence of abnormal contractions is indicative of arrest in the development of normal detrusor peroneal synergism, possibly due to chronic constipation.

Another possibility is that day wetting might induce voluntary peroneal contraction resulting in functional constipation. However, the elimination of enuresis and urinary tract infection after the treatment for constipation, as well as the fact that constipation and secondary encopresis preceded urinary symptomatology in several patients, militates against this possibility.

It is possible that the compression of the bladder due to pressure from a rectal fecal reservoir might serve to trigger such uninhibited bladder contractions. However, such contractions are not observed in pregnancy when the bladder is compressed by the gravid uterus. Since the innervation of the rectum and bladder are both from S2-4, it is possible that the development of constipation might precipitate, due to stimulation of the detrusor and consequent incoordination between detrusor contraction and external bladder sphincter relaxation.

One can surmise that in chronically severe constipated patients the rectum is never empty. In this case, the external sphincter of the anus remains the main if not the only means of rectal continence. It then is contracted continuously by the normal reflex mechanism and by voluntary action. Anal electromyography has been reported to be abnormal in children with urinary tract infection [Bailey et al. 1970]. Since concomitant contraction of the external sphincter of the bladder and the anus normally occurs, the presence of a continuous contraction of the anal

sphincter would induce the continuous contraction of the bladder sphincter resulting in urinary symptoms. This may explain the observation of Bailey et al. [1970] who noted a 57% incidence of abnormal anal electromyography in children with recurrent urinary tract infection. Elimination of constipation would result in cessation of voluntary anal and consequent urethral sphincter contraction, thus allowing for attainment of a more normal voiding pattern. This theory best explains the dramatic success of elimination of constipation in our patient series and is substantiated by the observation of an extremely high incidence of urinary tract infection in children with Hirschsprung's disease [O'Regan and Yazbeck 1984].

Pharmacologic manipulation has been utilized in the management of children with dysfunctional voiding problems with some success [Firlit et al. 1978, Smey et al. 1978]. However, pharmacologic manipulation based on urodynamic studies may result in variable symptomatic improvement without elimination of a possible primary cause i.e., functional constipation. Our experience indicates that the association of constipation and recurrent urinary tract infection is a frequent uncommonly recognized symptom complex. Rectal examination should be performed on all patients with recurrent urinary tract infection.

#### REFERENCES

- Allen TD, Bright TC, III 1978 Urodynamic patterns in children with dysfunctional voiding problems. *J. Urol.* 119: 247
- Bailey JA, Powers JJ, Waylonic GW 1970 A clinical evaluation of electromyography of the anal sphincter. *Arch. Phys. Med. Rehab.* 51: 403
- Berger RM, Maizels M, Moran GC, Conway JJ, Firlit CF 1983 Bladder capacity (ounces) equals age (years) plus 2 predicts normal bladder capacity and aids in diagnosis of abnormal voiding patterns. *J. Urol.* 129: 347
- Firlit CF, Smey P, King LR 1978 Micturition urodynamic flow studies in children. *J. Urol.* 119: 250
- Meunier P, Mollard P, Jaubert de Beaujeu M 1967 Manometric studies of anorectal disorders in infancy and childhood: an investigation of the physiopathology of continence and defecation. *Br. J. Surg.* 63: 402
- Neumann PZ, deDomenico IJ, Nogrady MB 1973 Constipation and urinary tract infection. *Pediatrics* 52: 241
- O'Regan S, Yazbeck S 1984 An increased incidence of urinary tract infection in children with Hirschsprung's disease. Abstract, Royal College of Physicians and Surgeons of Canada, Montreal, September 10-14
- Shopfner CE 1968 Urinary tract pathology associated with constipation. *Radiology* 90: 865
- Smey P, Firlit CF, King LR 1978 Voiding pattern abnormalities in normal children: Results of pharmacologic manipulation. *J. Urol.* 120: 574

**CONSTIPATION: A CAUSE OF ENURESIS, URINARY TRACT INFECTION AND VESICO-URETERAL REFLUX IN CHILDREN**

S. O'Regan and S. Yazbeck. Pediatrics and Surgery Department, Nephrology Division, Université de Montréal, Centre de Recherche Pédiatrique, Hôpital Sainte-Justine, 3175 Chemin Sainte-Catherine, Montreal, Quebec, Canada H3T 1C5. (reprint requests to SOR)

ABSTRACT

The observation that constipation alone may induce uninhibited bladder contractions in children and is associated with recurrent urinary tract infection in childhood suggests an etiologic association (1). Rectal distension due to faecal retention in chronic functional constipation causes bladder distortion (2) and may cause stimulation of detrusor stretch receptors resulting in detrusor peroneal dyssynergism. Distortion of the trigonal area may result in failure of ureteral valve competence and allow for vesico-ureteric reflux.

INTRODUCTION

The causes of recurrent urinary tract infection, enuresis and vesico-ureteric reflux in the absence of anatomic abnormality are unknown. Studies investigating recurrent urinary tract infection in children in the absence of radio-urollogic abnormalities have concentrated on the identification of particular characteristics of infecting bacteria resulting in urinary tract infection such as pili formation and host bladder wall receptors (3). The latter postulate infers that urinary tract infection is caused by infecting organism characteristics suggesting that it may be an infectious disease. This is not so. Enuresis has been attributed to abnormalities of neural control of bladder function (4) and has been treated with alarm systems (5) or with systemically acting drugs such as Imipramine and Vasopressin to decrease urinary output (7), none with a satisfactory result. Vesico-ureteric reflux in children has been attributed to abnormalities of ureteric insertion of the ureter into the bladder wall (8). However, the observations that reflux spontaneously resolves suggests that previous congenital anatomic abnormalities spontaneously resolve, an unlikely proposition (9).

Recurrent urinary tract infection in children

Recurrent urinary tract infection in children as shown by epidemiologic studies has an incidence of up to 3% in North American children (10).

Treatment usually consists of antibiotic therapy of infectious episodes or chronic prophylactic therapy. Urinary tract infection usually resolves at the end of the first decade or beginning of the second decade of life. In 1967, Shofpner et al. studying bladder and ureteral distortion due to constipation in 36 children noted that 3 of them had recurrent urinary tract infection (2). Neumann et al. noted the frequency of abnormal bowel habits, specifically constipation, in association with urinary tract infection in children and noted a high incidence of resolution of recurrent urinary tract infection with treatment of the constipation (11).

In a study of 47 children (1) it was noted, utilizing urodynamic and rectal manometric studies, that constipation was a constant associated condition. Constipation was measured not subjectively but objectively by measuring rectal ampulla capacity and rectal sphincter response to air balloon insufflation thus providing a measurement of constipation not subject to observer bias. Fifty percent of the mothers denied constipation as a symptom though questioning the individual children indicated defecation episodes of 2 to 3 times per week in the majority of cases. Aggressive treatment of the constipation resulted in resolution of all the recurrent urinary tract infections and also of associated enuresis in 67% of those who had it. These studies indicated the association of constipation with urinary tract infection.

A retrospective study of the incidence of urinary tract infection in Hirschsprung's disease (12), a constipating congenital abnormality of the large bowel, indicated an incidence greater than that normally observed in infancy. The study suggests that constipation may contribute to the development of recurrent urinary tract infection in children. The fact that the rectum may be abnormal in these children, substantiates the studies of Bailey et al. (13) who demonstrated abnormal anal electromyography in 57% of children with urinary tract infection.

Three possibilities exist as to how constipation may be associated with urinary tract infection:

- a) the presence of recurrent urinary tract infection may allow for the development of constipation;
- b) the presence of uninhabitable bladder contractions against a closed ureteral sphincter would lead to intermittent urinary wetting. Efforts to maintain continence would lead to ureteral and simultaneous anal sphincter contractions resulting in constipation;
- c) the presence of constipation with a faecal reservoir would compress the bladder causing stimulation of stretch receptors resulting in uninhibited bladder contractions. Detrusor contraction would force urine down into the urethra against the closed external sphincter resulting in reflux of urine contaminated with bacteria back up from the urethra resulting in bladder infection;
- d) chronic constipation is associated with external anal sphincter contractions (voluntary or reflex). This leads to simultaneous bladder external sphincter contraction leading to failure of normal development of detrusor contraction/peroneal relaxation synergism.

### Constipation and enuresis

Urodynamic studies in children with constipation in the absence of other symptomatology demonstrated uninhibited bladder contractions (1). Further studies in children with enuresis also demonstrated uninhibited bladder contractions with decreased bladder capacity as seen in children with recurrent urinary tract infection (1). Aggressive treatment of constipation in boys and girls with enuresis resulted in resolution of their enuresis. A decrease in bladder capacity is noted in enuretics (14) and may be due to bladder compression by the rectum. Shopfner (2) noted that 54% of 36 patients with constipation were enuretic. It was also noted by Bailey et al. (13) that 55% of the children with enuresis had abnormal anal electromyography substantiating the postulate indicating that constipation is an etiologic factor.

### Vesico-ureteric reflux and constipation

Several series of children with Hirschsprung's disease have shown mega-ureter with dilatation of the genito-urinary tract (15,16,17). Kottmeir et al. noted a similar incidence of this phenomenon in children with functional constipation (17). Also constipation has been observed to cause hydronephrosis (18,19). Surgical resolution of Hirschsprung's disease may result in return to normal of the uretero-calycial dilatation (20). Since innervation of the bladder in Hirschsprung's disease is normal (21), a similar neuro-etiology for bladder and ureteric dilatation is not tenable. Children with vesico-ureteric reflux in the absence of anatomic abnormalities have uninhibited bladder contractions with dysfunctional voiding patterns (22). Treatment with cholinergic drugs such as Oxybutyrin may result in a decrease in the degree or resolution of the reflux by depressing uninhibited bladder contractions (22). Our observation that children with vesico-ureteric reflux are commonly constipated suggested to us that a spectrum of abnormalities of recurrent urinary tract infection, enuresis and vesico-ureteric reflux all of which are associated with uninhibited bladder contractions in childhood may be secondary to bladder distortion by chronic functional constipation. Since aggressive treatment of constipation results in resolution of enuresis, and recurrent urinary tract infection, with resolution of non inhibited bladder contractions, we propose that vesico-ureteric reflux on which volumes have been written (20) may be caused by chronic constipation.

The constipation may be undetected because:

- a) physician does not inquire;
- b) the mother is unaware since she will not document bowel motions in children when it is not a primary complaint;
- c) because its presence may be over shadowed by symptomatology from recurrent urinary tract infection or psychologic trauma produced by enuresis;
- d) because of bowel preparation prior to radiologic investigation.

The cumulative evidence may be summarized as follows:

- 1) Uninhibited bladder contractions are common to enuresis, recurrent urinary tract infection and vesico-ureteral reflux.

- 2) Constipation is associated with recurrent urinary tract infection in children. Treatment of constipation results in resolution of infection.
- 3) Constipation is associated with obstructive uropathy and reflux in children with Hirschsprung's disease and functional megacolon. Urinary tract infection is increased in Hirschsprung's disease with or without obstructive uropathy. Treatment of the constipation results in resolution of urinary tract dilatation.
- 4) Enuresis and decreased bladder capacity is associated with constipation.
- 5) Posterior bladder wall distortion may be caused by constipation.
- 6) Constipation may be the common etiologic factor causing bladder distortion and uninhibited detrusor contraction resulting in enuresis, recurrent urinary tract infection and vesico-ureteral reflux.

#### REFERENCES

1. O'Regan S, Yazbeck S, Schick E. Constipation unstable bladder, urinary tract infection syndrome. Submitted for publication.
2. Shopfner CE. Urinary tract pathology associated with constipation. *Radiology* 90: 865, 1968.
3. Swenson SB, Hultberg H, Källenius G, Lorchonen TK, Möllby R, Winberg J. P-fimbriae of pyelonephritogenic E. Coli: identification and chemical characterization of receptors. *Infection* 11: 61, 1983.
4. Whiteside CG, Arnold EP. Persistent primary enuresis; a urodynamic assessment. *Brit Med J* 1: 364, 1975.
5. Doleys DM. Behavioral treatments for nocturnal enuresis in children: a review of the recent literature. *Psychol Bull* 84: 30, 1977.
6. Kass EJ, Diokno AC, Montealegro A. Enuresis: principles of management and results of treatment. *J Urol* 121: 794, 1979.
7. Terho P, Kekomaki M. Management of nocturnal enuresis with a vasopressin analogue. *J Urol* 131: 925, 1984.
8. Stephens FD. Urologic aspects of recurrent urinary tract infection in children. *J Pediatr* 80: 725, 1972.
9. Normand C, Smellie J. Vesico-ureteral reflux: the case for conservative management. in *Reflux Nephropathy*. (J Hodson, P Kincaid-Smith, eds) Masson, New York, 1979.
10. Kunin CM, Deutscher R, Paquin A. Urinary tract infections in school children: epidemiologic, clinical and laboratory study. *Medicine* 30: 91, 1964.
11. Neumann PZ, De Domenico IJ, Nogrady MB. Constipation and urinary tract infection. *Pediatrics* 52: 241, 1973.
12. O'Regan S, Yazbeck S. An increased incidence of urinary tract infection in children with Hirschsprung's disease. Abstract Royal College of Physicians and Surgeons of Canada, Montreal, Sept 10-14, 1984.
13. Bailey JA, Powers JJ, Waylonis GW. A clinical evaluation of electromyography of the anal sphincter. *Arch Phys Med Rehab* 51: 403, 1970.

14. Berger RM, Maizels M, Moran GC, Conway JJ, Firlit CF. Bladder capacity (ounces) equals age (years) plus 2 predicts normal bladder capacity and aids in diagnosis of abnormal voiding patterns. *J Urol* 129: 347, 1983.
15. Ehrenpreis T. Hirschsprung's disease. Year Book Medical Publishers, Chicago, p 60, 1970.
16. Swenson O, MacMahon HE, Jacques WE, Campbell JS. New concept of etiology of megalo-ureters. *New Engl J Med* 246: 41, 1952.
17. Kottmeier PK, Clatworthy DW. Aganglionic and functional megacolon in children - a diagnostic dilemma. *Pediatrics* 36: 572, 1965.
18. Ravich L, Lerman PH, Schell NB. Urinary retention due to fecal impaction. *NY J Med* 63: 3289, 1963.
19. Savage JP. The deleterious effect of constipation upon the re-implanted ureter. *J Urol* 109: 501, 1973.
20. Sieber WK, Soave F. Hirschsprung's disease. Current problems in surgery. *Surgery* 15: 14, 1978.
21. Lebowitz S, Bodian M. A study of the vesical ganglia in children and the relationship to the megaureter-megacystis syndrome and Hirschsprung's disease. *J Clin Pathol* 167: 342, 1963.
22. Koff SA, Murtagh DS. The uninhibited bladder in children: effect of treatment of recurrence of urinary tract infection and on vesico-ureteral reflux resolution. *J Urol* 130: 1138, 1983.



## Constipation and the Urinary System

A major drawback in studying relationships between constipation and abnormalities of the urinary system has been the lack of a standard definition of constipation. In addition, a method of measurement of rectal abnormalities which could be correlated with the presence of constipation and which was recordable has only recently been available. The availability of rectal manometry has allowed for the assessment of rectal function in evaluation of constipation. We consider constipation to be present if one or more of the following criteria are present:

- More than 72 h interval between bowel movements.
- The presence of overflow fecal incontinence (encopresis or soiling).
- The passage of small hard scybalous stools with intermittent large stools.
- Incomplete rectal emptying as determined by rectal examination after defecation.
- Grossly decreased level of perception and increased tolerance to balloon insufflation in the presence of a normal rectoanal relaxation on rectal manometry, combined with any of the above elements.<sup>1-4</sup>

Rectal manometry allows the operator to determine the smallest rectal distention perceived by the patient, the volume at which relaxation of the internal sphincter occurs and the maximal volume that can be tolerated by the patient without discomfort or pain.<sup>1</sup>

This facilitates the diagnosis of constipation, especially as norms for manometry have already been established for children. The diagnosis of constipation is to be made by a physician with appropriate questioning on quantity and quality of stools and frequency of defecation, and physical examination. A simple question to the mother as to whether the child is constipated or not allows for a misdiagnosis to be made on the basis of arbitrary definitions. A convenient way of avoiding being misled is to obtain a written record of stool frequency and quality at follow up visits.

Similarly, the advent of techniques to assess urodynamics has led to the identification of non-inhibited contractions<sup>2</sup> of the bladder in patients with urinary tract infection, and on occasion enuresis and vesicoureteral reflux.

### Constipation and Urinary Obstruction

Prior to puberty the child's abdominal and pelvic cavities are one. As a result the close proximity of the rectum to the

posterior wall of the bladder is such that any gross distension of the rectum by stools has been reported to result in compression of the bladder, with bladder neck obstruction or distension of the urethra leading to urinary obstruction.<sup>3</sup>

### Constipation and Urinary Tract Infection

Leishman, in 1939, suggested that constipation did not play a role in urinary tract infection in adults.<sup>8</sup> In a series of 39 patients, Shopfner demonstrated, in 1968, that the presence of constipation has the potential for distorting the bladder wall and urethra.<sup>9</sup> Of the 39 patients, eight had reflux, 21 had enuresis and two had urinary tract infection. Campbell<sup>10</sup> described the lazy bladder syndrome as consisting of recurrent cystitis, urinary dribbling and constipation. Neumann *et al.*<sup>11</sup> in 1973, noted the presence of constipation in children who presented with anomalies ranging from recurrent urinary tract infection to mild or severe vesicoureteral reflux. Constipation was diagnosed on the basis of a history of infrequent bowel motions and the radiologic evidence of accumulated feces. Aggressive treatment of constipation resulted in resolution soiling, with proven rectal reservoirs of feces as determined by rectal examination and rectal manometry. It is essential therefore that a written record of stool frequency and quality be kept so that parents recognize the presence of constipation. This then facilitates compliance with appropriate therapy.

An increased incidence of bacteriuria occurs in rats in whom fecal retention is surgically induced.<sup>17</sup>

### Constipation and Enuresis

The etiology of enuresis is not completely understood. In 17 constipated enuretics who had uninhibited detrusor contractions, treatment of coexisting constipation resulted in the cessation or improvement of enuresis in 15 and two cases respectively.<sup>18</sup>

As previously stated, Shopfner, in discussing urinary tract pathology associated with constipation, also noted that 54% of these patients were enuretic.<sup>9</sup> Baumann and Hinman<sup>19</sup> discussed the treatment of incontinence with non-obstructive disease of the urinary tract. Their 73 male patients had enuresis with encopresis. Though the emphasis was on hypnotherapy as specific treatment of enuresis, they did note that aggressive treatment of constipation, commonly including digital evacu-

ation of the rectum, was required before cessation of enuresis, suggesting that constipation may have been a relevant element causing enuresis. Urodynamics in patients who were severely constipated but did not have urinary symptoms showed uninhibited bladder contractions.<sup>18</sup> This suggests that constipation may induce uninhibited bladder contractions, with consequent enuresis.

### Constipation and Vesicoureteral Reflux

Dilatation of the urinary tract, including the bladder, has been noted by numerous authors to occur in association with Hirschsprung's disease.<sup>20-21</sup> Successful treatment of this disease may result in resolution of the vesicoureteral reflux.<sup>22</sup> Kottmeier and Clatworthy noted a higher incidence of vesicoureteral reflux in children with severe functional constipation than in those with Hirschsprung's disease.<sup>23</sup> Also, constipation may cause ureteral dilatation with hydronephrosis and, when relieved, may allow for resolution of urinary tract dilatation.<sup>24</sup> Ochoa and Gorlin have described a syndrome of distortion of facial expression associated with urinary tract dilatation.<sup>25</sup> In two-thirds of afflicted patients moderate to severe constipation was present. Whether the constipation is a primary motility problem contributing to urinary tract dilatation or is secondary to chronic dehydration due to renal failure associated polyuria was not determined. Abnormal large bowel motility has been noted in patients with vesicoureteral reflux.<sup>26-27</sup>

We have noted the presence of constipation in children with primary vesicoureteral reflux.<sup>28</sup> Although a control study was not done, rapid resolution of reflux occurred with aggressive treatment of constipation in the absence of antibiotic or anticholinergic therapy for treatment of associated uninhibited bladder contractions. White also noted that resolution of infection with reflux was more easily attained when associated constipation was aggressively treated.<sup>29</sup> Hinman<sup>30</sup> observed that the constipated child may have an evolution of encopresis to enuresis to urinary tract infection to vesicoureteral reflux. Studies by Bailey *et al.*<sup>31</sup> noted a 55% incidence of abnormal anal sphincter electromyograms in children with enuresis and urinary tract infection, suggesting the possibility that abnormalities of the rectum may cause enuresis and urinary tract infection and reflux in children. Indeed the observation that children with functional constipation may have uninhibited contractions of the bladder in the absence of urinary symptoms strongly supports this possibility.

Because the external anal sphincter and the urethral sphincter together with the perineal musculature may be considered to be a single physiologic unit, with voluntary anal contraction consequent concomitant urethral sphincter contraction occurs. It is possible that the urethral sphincter contraction may be responsible for a dyssynergic voiding pattern with secondary bladder instability, enuresis, urinary tract infection or vesicoureteral reflux.

Since the rectum may be dilated for months to years in association with chronic constipation, one-time emptying of the rectal ampulla will result in a momentarily empty but still

dilated rectum, without resolution of constipation. Oral therapy alone directed to the maintenance of an empty rectum is not consistently successful; the maintenance of an empty rectum is only achieved by repeated use of enemas. A satisfactory approach is the use of daily saline enemas (5 ml of salt in 1 liter of warm water). Using an enema bag held approximately 0.5 m above the patient maintained in a proper position, an adequate volume will be infused under gravity with minimal induction of cramps.

Although the frequency of enemas will be decreased progressively, treatment of constipation should be continued until rectal tone returns to normal. This usually takes at least 3 months.

### Summary


The morbidity associated with enuresis, urinary tract infection and vesicoureteral reflux is exceedingly high. Constipation is a condition easily amenable to resolution with appropriate therapy. Consequently, in the management of children with any of these conditions, appropriate investigation should be performed to determine the presence of constipation. Should constipation be present, therapy should be instituted so that it may aid in the resolution of the urinary symptoms.

### References

1. Meunier P., Mollard P. and Jaubert de Beaujeu M. Manometric studies of anorectal disorders in infancy and childhood: an investigation of the physiopathology of continence and defecation. *Br. J. Surg.* 63, 402 (1967).
2. Abrams P., Blauas J.G., Stanton S.L. and Andersen J.T. The standardisation of terminology of lower urinary tract function. *Scand. J. Urol. Nephrol. Suppl.* 114 (1988).
3. Jewett H.J. Diagnosis and management of urinary tract obstructions in children. *Surg. Clin. N. Am.* 32, 1371 (1951).
4. Grunberg A. Acute urinary obstruction due to fecal impaction. *J. Urol.* 83, 301-302 (1960).
5. Ravich L., Lerman P.H. and Schell N.B. Urinary retention due to fecal impaction. *N.Y. J. Med.* 63, 3289-3291 (1963).
6. Gallo R. and Presman D. Urinary retention due to fecal impaction in children. *Pediatrics* 45, 292-294 (1970).
7. Shandling B. and Desjardins J.G. Anal myomectomy for constipation. *J. Pediatr. Surg.* 4, 1 (1969).
8. Leishman W.D. and Oxford D.M. Bacillus coli infection of the urinary tract. *Lancet* ii, 971-973 (1939).
9. Shopfner C.E. Urinary tract pathology associated with constipation. *Radiology* 90, 865-877 (1968).
10. Campbell N.A. Functional abnormalities of the bladder in children: characteristics of chronic cystitis and chronic urethritis. *J. Urol.* 104, 926-929 (1970).
11. Neumann P.Z., de Dominico I.J. and Nogrady M.B. Constipation and urinary tract infection. *Pediatrics* 52, 241-245 (1973).
12. White R.H.R. and Taylor C.M. The non-operative management of primary vesicoureteric reflux. In *Management of Vesicoureteric Reflux* (Johnston ed.), pp. 117-136. Baltimore: Williams and Wilkins (1984).
13. Smellie J.M., Katz G. and Grunbert R.N. Controlled trial of prophylactic treatment in childhood urinary tract infection. *Lancet* ii, 175-178 (1978).
14. Yazbeck S. and O'Regan S. Hirschsprung's disease and urinary tract infection: unrecognized association. *Nephron* 43, 211-213 (1986).
15. Vargas J.H., Sachs P. and Ament M.E. Chronic intestinal pseudo-obstruction syndrome in pediatrics. Results of a national survey by members of the North American Society of Pediatric gastroenterology and nutrition. *J. Pediatr. Gastroenterol. Nutr.* 7, 323-332 (1988).
16. O'Regan S., Yazbeck S. and Schick E. Constipation, bladder instability, urinary tract infection syndrome. *Clin. Nephrol.* 23 152-154 (1985).

17. Breda G., Bianchi G.P., Bonimi U., Piacentini I. and Farello G. Faecal stasis and bacteriuria: experimental research in rats. *Urol. Res.* 2, 155-157 (1975).
18. O'Regan S., Yazbeck S., Hamberger B. and Schick E. Constipation a commonly unrecognized cause of enuresis. *Am. J. Dis. Child.* 140, 260-261 (1986).
19. Baumann F.W. and Hinman F. Treatment of incontinent boys with nonobstructive disease. *J. Urol.* 111, 114-116 (1974).
20. Ehrenpreis T. *Hirschsprung's Disease*, pp. 60-61, Chicago: Yearbook Medical Publishers (1970)
21. Swenson O., MacMahon H.E., Jacques W.E. and Campbell J.S. New concept of etiology of megalo-ureters. *N. Engl. J. Med.* 476, 41-46 (1952).
22. Sieber W.K. and Soave F. Hirschsprung's disease. *Curr. Probl. Surg.* 15, 14-15 (1978).
23. Kottmeier P.K. and Clattworthy D.W. Aganglionic and functional megacolon in children - a diagnostic dilemma. *Pediatrics* 36, 572-582 (1965).
24. Savage J.P. The deleterious effect of constipation upon the reimplanted ureter. *J. Urol.* 109, 501-503 (1973).
25. Ochoa B. and Gorlin R.J. Orofacial (ochoa) syndrome. *Am. J. Med. Genet.* 27, 661-667 (1987).
26. Salov P.P. Vesico-ureteral reflux associated with functional disorders of the distal segment of the large intestine in children. *Urol. Nefrol. (Mosk)*. 17-22 (1988).
27. Baldew I.M. Children with urination problems in constipation. *Med. Tijdschn. Geneesk.* 129, 3-5 (1985).
28. O'Regan S., Schick E., Hamberger B. and Yazbeck S. Constipation associated with vesico-ureteral reflux. *Urology* 28, 394-396 (1986).
29. White R. Reflux nephropathy. Update 1983. Discussion. *Contr. Nephrol.* p. 249 (Karger, Basel 1984).
30. Hinman F. Urinary tract damage in children who wet. *Pediatrics* 54, 143-150 (1974).
31. Bailey J.A., Powers J.J. and Wayeonis G.W. A clinical evaluation of electromyography of the anal sphincter. *Arch. Phys. Med. Rehabil.* 51, 403-408 (1970).

# Daily Enema Regimen Is Superior to Traditional Therapies for Nonneurogenic Pediatric Overactive Bladder

Global Pediatric Health  
Volume 3: 1–4  
© The Author(s) 2016  
Reprints and permissions:  
DOI: 10.1177/2333794X16632941  
gph.sagepub.com  


Steve J. Hodges, MD<sup>1</sup>, and Marc Colaco, MD<sup>1</sup>

## Abstract

Our objective was to evaluate the efficacy of daily enemas for the treatment of overactive bladder (OAB) in children. This study was a prospective, controlled trial of 60 children with nonneurogenic OAB. The control patients (40) were treated with standard therapies, including timed voiding, constipation treatment with osmotic laxatives, anticholinergics, and biofeedback physical therapy, whereas the treatment patients (20) received only daily enemas and osmotic laxatives. On assessment of improvement of OAB symptoms, only 30% of the traditionally treated patients' parents reported resolution of symptoms at 3 months, whereas 85% of enema patients did. At the onset of the study, the average pediatric voiding dysfunction score of all patients was 14, whereas on follow-up, the average scores for traditionally treated patients and enema-treated patients were 12 and 4, respectively. This study demonstrated that daily enema therapy is superior to traditional methods for the treatment of OAB.

## Keywords

voiding dysfunction, dysfunctional elimination, constipation, enema, incontinence

Received December 29, 2015. Received revised December 29, 2015. Accepted for publication January 12, 2016.

## Introduction

Overactive bladder (OAB) is a common and vexing problem in children, and despite great advancement in therapies, a certain percentage of patients remains resistant to treatment. We believe that children whose symptoms do not resolve with timed voiding, laxatives, anticholinergic medications, and biofeedback physical therapy do so because of an undiagnosed and inadequately treated megarectum and that therapy directed specifically at the dilated rectum will resolve OAB symptoms most efficaciously. In this study, we evaluated the efficacy of daily enemas for the treatment of OAB in children.

## Material and Methods

The study was approved by the institutional review board. This was a prospective, controlled trial of 60 children with nonneurogenic OAB. The inclusion criterion was a diagnosis of pediatric nonneurogenic OAB. Exclusion criteria included neurogenic cause of bladder dysfunction, urinary tract infection, prior lower-urinary-tract surgery, and any diagnosed anatomical abnormalities of the urinary tract that could influence voiding function, such as posterior urethral valves. OAB was

defined as uncontrolled daytime urge incontinence, and bladder function was measured using the pediatric voiding dysfunction symptom score (DVSS).

The 40 control patients were treated with traditional therapies, including timed voiding, osmotic laxative PEG3350 (regardless of bowel history to maintain daily, soft bowel movements), and in select cases, anticholinergic medications and/or biofeedback therapy. The 20 remaining patients were prescribed only a daily enema (liquid glycerin suppository for ages 2 to 5, pediatric fleet enema for ages 6 to 11) and enough osmotic laxative to maintain soft spontaneous bowel movements, with no other therapy or voiding schedule. If the voiding symptoms resolved while on the daily enemas, patients were instructed to taper off the daily enemas over a 2-month time period (an enema every other day for a month, and then an enema twice weekly for a month). All patients were evaluated on each visit with complete

<sup>1</sup>Wake Forest University School of Medicine, Winston Salem, NC, USA

### Corresponding Author:

Steve J. Hodges, Wake Forest University School of Medicine, Medical Center Boulevard, Winston Salem, NC 27157, USA.  
Email: shodges@wakehealth.edu



Creative Commons CC-BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 3.0 License (<http://www.creativecommons.org/licenses/by-nc/3.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access page (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

**Table 1.** Pretreatment Measurements.<sup>a</sup>

	Treatment Group	Control Group	Difference (P)
DVSS	14	14	0 (.68)
BSS	3.55	3.95	0.4 (.12)
Rome III	0.55	0.85	0.3 (.3)
Rectal diameter (cm)	6.35	6.2	0.15 (.66)

Abbreviations: DVSS, pediatric voiding dysfunction symptom score; BSS, Bristol Stool Scale.

<sup>a</sup>There were no variables that were significantly different between the control and treatment groups prior to enema.

**Table 2.** Measurements Done After the Treatment Phase.<sup>a</sup>

	Treatment Group	Control Group	Difference (P)
DVSS	4	12	
Change from pretreatment	-10	-2	8 (<.01)
BSS	3.85	4.05	
Change from pretreatment	0.3	0.1	0.2 (.08)
Rome III	0.1	0.2	
Change from pretreatment	0.45	0.65	0.2 (.45)
Rectal diameter (cm)	2.15	5	
Change from pretreatment	4.2	1.2	3 (<.01)

Abbreviations: DVSS, pediatric voiding dysfunction symptom score; BSS, Bristol Stool Scale.


<sup>a</sup>Participants who underwent enemas had a significantly greater improvement in DVSSs and maximum rectal diameters.

history and physical, urinalysis, Bristol Stool Scale (BSS), Rome III criteria, KUB X-ray, and Pediatric Voiding Dysfunction Questionnaire. All children were followed up at 3 months.

Data analysis was performed using SPSS Statistics Version 23 (IBM Corp, Armonk, NY). For the nonparametric variables—DVSSs, BSS scores, and Rome III scores—comparisons were made using Mann-Whitney *U* tests. For the continuous variable—maximum rectal diameter on KUB—a student *t* test was used. Comparisons were made both within groups for the pretreatment and posttreatment phases as well as between groups.

## Results

A total of 60 children (20 experimental and 40 controls) were included in this study. Table 1 demonstrates the mean DVSS, BSS, Rome III, and rectal diameters prior to treatment. There was no significant difference on any of these measurements between the control and treatment groups. Table 2 demonstrates the mean posttreatment measures and the mean change in each metric after the treatment period. Patients who underwent enema had significantly more improvement in DVSSs




**Figure 1.** Pretreatment and posttreatment pediatric voiding dysfunction symptom score (DVSSs): the treatment group showed a significantly greater improvement in DVSS when compared with the control group.

and significantly greater change in maximum rectal diameters than control patients (Figures 1 and 2). There was no difference in the amount of change between groups. Of note, both control and treatment groups demonstrated significant improvement in all measured variables after the treatment period.

## Discussion

The historical teaching regarding OAB of childhood was that a congenital obstruction interrupted urine flow and led to the development of detrusor hypertrophy and hyperactivity, and the accepted therapy was serial and repeated dilation of this obstruction.<sup>1</sup> As time progressed, this anatomical obstruction was found to be the result of a willful dyssynergic contraction of the pelvic floor during voiding, and the treatment was changed to biofeedback physical therapy.<sup>2</sup> This nonphysiological contraction of the urethral sphincter was no medical curiosity but a disease process so severe that it could influence the natural history of many childhood disorders, such as vesicoureteral reflux and nocturnal enuresis, and in extreme cases induce renal failure.<sup>3,4</sup>

There is some debate among scientists as to whether this dyssynergic sphincter contraction is a learned or inborn condition.<sup>5</sup> It is the author's opinion that in children with an intact nervous system, uninhibited voiding dominates the infantile period of voiding prior to toilet training. This is most clearly represented by the progressive bladder growth in pre-toilet-trained



**Figure 2.** Pretreatment and posttreatment maximum rectal diameters: the treatment group also showed a significantly greater improvement in maximum rectal diameter on KUB when compared with the control group.

children, with increasing compliance.<sup>6</sup> This is mirrored in children with cerebral palsy, who demonstrate progressive bladder growth when they maintain an uninhibited, infantile voiding pattern.<sup>7</sup>

The teaching of bladder overactivity as the natural progression of an obstructed voiding pattern evolved from pathological models of voiding dysfunction such as posterior urethral valves and the neurogenic bladder, which develops in myelodysplastic patients with discoordinated sphincters.<sup>8-10</sup> This led to the development of the modern model of dysfunctional elimination, an acquired condition where children paradoxically fail to relax their pelvic floor during elimination, resulting in bowel and bladder pathology.<sup>11,12</sup> This study sought to investigate an alternative theory regarding the origins of dysfunctional elimination, with a therapy directed toward that cause to examine its benefits.

We have long known of the association between bowel and bladder dysfunction. This was first noticed in the 1960s, largely because of the presence of urinary symptoms in children with Hirschsprung's disease. In his seminal work, Shopfner<sup>13</sup> noted that the distention of the colon, especially the rectum, could have profound effects on bladder function.

O'Regan furthered this work with several groundbreaking studies linking rectal distention to bladder overactivity, with excellent success in treating nocturnal enuresis, urinary tract infections, and vesicoureteral reflux simply by alleviating this rectal distention. What has unfortunately hampered the development of this

work has been the lack of a uniform definition of constipation.<sup>14-17</sup>

O'Regan defined constipation not as functional constipation, but mainly as the presence of fecal soiling, incomplete rectal emptying, and/or grossly decreased level of perception to balloon insufflation on anorectal manometry. The interesting discovery that O'Regan made was that often children with OAB symptoms presented with no functional signs of constipation, yet had markedly abnormal anorectal manometry studies. In other words, they often volitionally delayed defecation until the rectum distended to fill the anatomical pelvis and then would reach a new, abnormal homeostasis where stools would evacuate at regular intervals and with surprisingly normal appearance; yet the rectal tone would be so diminished as to have abnormal manometry studies. Put another way, these children were changing the rectum from a sensing organ (which in normal circumstances provides cues on the need to defecate), to a storage organ, with decreased sensation, and often resultant fecal soiling, but more often than not—normal stooling patterns.<sup>16</sup>

O'Regan early on discovered what we have also demonstrated in this study that parental reporting of their children's bowel habits by BSS or Rome III criteria is often inaccurate, and even when accurate, often not helpful in diagnosing rectal distention in children with OABs.<sup>17</sup> Not only that, but in children with completely normal bowel habits, rectal distention can be the main or sole cause of urinary symptoms, leading to compression of the bladder, uninhibited contractions, and often urethral obstruction, all of which has been proved years ago.<sup>13</sup>

The ability of rectal stool to induce uninhibited bladder contractions is well understood and was described as early as the 1980s; this work has been supported by numerous studies, although the exact mechanisms have yet to be defined. The uncanny ability of children to be cured of nocturnal enuresis by simply restoring normal rectal tone is a great testament to this relationship.<sup>16</sup>

What O'Regan proposed was that “in chronically constipated children; the rectum is never empty, necessitating the repetition or maintenance of rectal sphincter complex contraction to maintain fecal continence. Consequent concomitant urethral sphincter contraction occurs”<sup>16</sup>p. 261. In other words, the pelvic floor contractions during voiding are not a willful process that can be unlearned, but a physiological response to stool withholding. And this could be reversed by directing therapy specifically at rectal dilation, in other words daily enemas, with excellent results.

So what if the modern theory of dysfunctional elimination and the resultant OAB is wrong? Our research

points to a different cause altogether. We have proved that simply emptying the rectum repeatedly with the goal of restoring normal rectal tone resolves OAB in children more efficaciously than the standard of care. In fact, if the current model of voiding dysfunction were accurate, it should be impossible for our treatment to have been beneficial at all because we made no effort to influence pelvic floor function. Some would argue that our therapy would make pelvic floor contractions worse. Yet, in almost all children, the bladder symptoms resolved. And in children whose symptoms did not resolve, the enemas were often not successful in restoring rectal tone.

## Conclusion

A daily enema regimen specifically targeted at restoring normal rectal tone is more effective than the standard of care for the treatment of OAB in children.

## Author Contributions

SJH contributed to the conception and design; contributed to the acquisition, analysis, and interpretation of data; drafted the manuscript; gave final approval; and agrees to be accountable for all aspects of work ensuring integrity and accuracy. MC contributed to the analysis and interpretation of data and drafted the manuscript.

## Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

## References

- Milanovic D, Sremcevic D, Perovic S, Scepanovic D, Krstic Z. Stenosis of the external urethral opening as one of the causes of recurrent urinary infections in girls: surgical treatment [in Serbian]. *Acta Chir Jugosl.* 1989;36:229-237.
- Palmer LS. Biofeedback in the management of urinary continence in children. *Curr Urol Rep.* 2010;11:122-127.
- Hinman F Jr. Nonneurogenic neurogenic bladder (the Hinman syndrome): 15 years later. *J Urol.* 1986;136:769-777.
- Sillen U. Bladder dysfunction and vesicoureteral reflux. *Adv Urol.* 2008;(2008):815472.
- Yeung CK. The normal infant bladder. *Scand J Urol Nephrol Suppl.* 1995;173:19-23.
- Zerin JM, Chen E, Ritchey ML, Bloom DA. Bladder capacity as measured at voiding cystourethrography in children: relationship to toilet training and frequency of micturition. *Radiology.* 1993;187:803-806.
- Richardson I, Palmer LS. Clinical and urodynamic spectrum of bladder function in cerebral palsy. *J Urol.* 2009;182:1945-1948.
- Hodges SJ, Patel B, McLorie G, Atala A. Posterior urethral valves. *ScientificWorldJournal.* 2009;9:1119-1126.
- Chung DE, Sandhu JS. Overactive bladder and outlet obstruction in men. *Curr Urol Rep.* 2011;12:77-85.
- Madersbacher H. Neurogenic bladder dysfunction in patients with myelomeningocele. *Curr Opin Urol.* 2002;12:469-472.
- Franco I. New ideas in the cause of bladder dysfunction in children. *Curr Opin Urol.* 2011;21:334-338.
- Koff SA. Relationship between dysfunctional voiding and reflux. *J Urol.* 1992;148:1703-1705.
- Shopfner CE. Urinary tract pathology associated with constipation. *Radiology.* 1968;90:865-877.
- O'Regan S, Schick E, Hamburger B, Yazbeck S. Constipation associated with vesicoureteral reflux. *Urology.* 1986;28:394-396.
- O'Regan S, Yazbeck S. Constipation: a cause of enuresis, urinary tract infection and vesico-ureteral reflux in children. *Med Hypotheses.* 1985;17:409-413.
- O'Regan S, Yazbeck S, Hamberger B, Schick E. Constipation a commonly unrecognized cause of enuresis. *Am J Dis Child.* 1986;140:260-261.
- O'Regan S, Yazbeck S, Schick E. Constipation, bladder instability, urinary tract infection syndrome. *Clin Nephrol.* 1985;23:152-154.

# Systematic review: the adverse effects of sodium phosphate enema

J. MENDOZA, J. LEGIDO, S. RUBIO & J. P. GISBERT

Department of Gastroenterology,  
Hospital Universitario de La Princesa,  
Madrid, Spain

Correspondence to:  
Dr J. P. Gisbert, Playa de Mojacar 29,  
Urb, Bonanza, 28669 Boadilla del  
Monte, Madrid, Spain.  
E-mail: gisbert@meditex.es

#### Publication data

Submitted 21 March 2007  
First decision 13 April 2007  
Resubmitted 18 April 2007  
Accepted 18 April 2007

## SUMMARY

### Background

Sodium-phosphate enemas are widely used to treat constipation, and are rarely associated with side effects.

### Aim

A systematic review of the literature was conducted to identify the most common adverse effects of sodium-phosphate enemas and associated risk factors.

### Methods

A systematic search was conducted in Internet (MEDLINE), and the Cochrane Library, from January 1957 to March 2007.

### Results

A total of 761 references were identified initially, and 39 relevant papers were finally selected. The most common therapeutic indications included constipation (63%). Sixty-eight per cent of the patients having adverse effects had associated conditions, the most common being gastrointestinal motility disorders, cardiological diseases and renal failure. Virtually, all side effects were due to water and electrolyte disturbances. Most patients were under 18 years of age (66%) or older than 65 years (25%). A total of 12 deaths were found.

### Conclusion

The main side effects caused by sodium phosphate enemas are water and electrolyte disturbances. The main risk factors are extreme age and associated comorbidity.

*Aliment Pharmacol Ther* 26, 9–20



## INTRODUCTION

Monosodium or disodium phosphate enemas are used for the treatment of acute and chronic constipation, and also for colon cleaning as preparation for endoscopic and surgical procedures, in both children and adults.<sup>1-3</sup> Phosphate enemas contain sodium acid phosphate and sodium phosphate, which have an osmotic activity. This activity could increase the water content and the volume of the stool, which will follow to a rectal distension. It is thought that this induces defecation with stimulation of rectal motility. Generally, the effect is limited to 5–10 min, which lowers the effect of phosphate toxicity as it is evacuated with the stool. These products have been widely used for many years, and have been associated with minimal adverse effects in the general population. However, there are reports in the literature of some clinical cases with severe side effects, even leading to death. If defecation does not take place, pooling of the fluid in the bowel can result in large amounts of water in the gut, causing dehydration. On the other hand, if phosphate is retained in the gut lumen can potentially be absorbed, and sudden and severe hypernatraemia and hyperphosphataemia may result.<sup>4</sup>

Manufacturers propose a careful use of the product in young children (<2 years) and in the elderly population, specially if associated comorbidity as renal disease or impaired intestinal motility exists. Nevertheless, there is a lack of information of the real risk of phosphate enemas. We therefore considered it necessary to conduct a systematic review to know what are exactly the most common side effects, their frequency, their severity and the profile of patients with a high risk of experiencing such complications.

## METHODS

We performed a literature search in Internet in the MEDLINE database (from January 1957 to March 2007). The clinical trials register (Cochrane Controlled Trials Register) of the Cochrane Library (number 1, 2007) was also reviewed. The following descriptors or key words were used (in all search fields): 'phosphate enema or sodium phosphate enema' or 'phosphate-based enema' or (phosphate AND enema) or (fleet AND enema) or 'sodium phosphate laxatives' or 'sodium phosphate catharsis' or 'sodium phosphate cathartic'. No restriction by language or by type of publication

was introduced. Literature references included in the papers meeting the selection criteria were also reviewed.

We selected the articles referring to secondary effects because of the administration of phosphate-based enemas. Data from the articles about anorectal or traumatic injuries were not examined because of the different ethiopathological approach.

To analyse the results, we performed a subanalysis according to the age criteria of the manufacturer's recommendations (children under 2 years, children under 18 years, adults and elderly (above 65 years), to perform a more comprehensive analysis.

In articles evaluating the side effects of sodium phosphate enemas, data were collected on the number of patients, sex, age, comorbidity, indication for use, number of units administered, deaths and their cause when they were due to use of enemas. Data extraction was conducted by two independent reviewers and discrepancies in the interpretation were resolved by consensus.

## RESULTS

The search conducted initially identified a total of 773 literature references. After a first selection by reading their abstracts, 707 references were discarded; most of these (553) did not refer to the question in hand. A further large group (146 references) was discarded because they addressed about sodium phosphate cleaning solutions administered by the oral route. Finally, 20 articles on sodium phosphate enemas were not included because their side effects were not reported.<sup>5-23</sup>

The remaining 54 articles were comprehensively analysed. Eleven of these were clinical trials, and seven of such trials compared several cleaning methods for performing endoscopy (sodium phosphate enemas, oral laxative sachets) and secondarily analysed adverse effects.<sup>2, 3, 24-27</sup> The remaining four clinical trials evaluated water and electrolyte disturbances after enema administration.<sup>28-31</sup> Of the remaining 43 references, we found conclusive data in 39 references, all of them case reports and letters to the editor.<sup>32-70</sup> We also found four references to enema-induced anorectal injuries.<sup>71-74</sup> All the cases revealed a damage in the anorectal tissue, mainly because of a harmful application. Because of the differential cause of damage, these cases were not considered as adverse effects

but as a consequence of a harmful application and they were not considered in the review.

Out of these 39 references,<sup>32–70</sup> the occurrence of side effects after administration of sodium phosphate enemas was reported in 44 patients. Twenty-two of these patients (50%) were women and 22 men (50%). Mean age of patients was 26 years (range, 6 weeks to 96 years).

### Therapeutic indications

Therapeutic indications included constipation in 28 patients (64%)<sup>33, 37–39, 41, 42, 44–50, 52, 54–70</sup> and preparation for diagnostic test (barium enema or colonoscopy) or surgery in six patients (14%).<sup>32, 36, 39, 57, 64, 69</sup> No mention was made of the indication in all other cases (10 patients, 23%).<sup>34, 35, 38, 40, 43, 51, 53</sup>

### Units administered

Units administered were difficult to assess, as dosage was not given for all patients. Dosage was stated in a total of 40 cases<sup>32, 33, 35–44, 46–56, 58–70</sup> (91%), but many of these data were incomplete or inadequate (not exact dosage, type of enema or frequency of administration). The exact dosage and frequency when it was given are in Tables 1–5. The maximum number of enemas received by a patient was 8,<sup>62</sup> but the time interval over which they were administered was not stated. The maximum frequency found in all analysed studies was six enemas over 6 h.<sup>69</sup>

### Comorbidity

Regarding past medical history, 38 patients (86%) had prior diseases,<sup>33, 36–44, 46, 47, 49–70</sup> as summarized below. The most common associated diseases were gastrointestinal conditions, found in a total of 18 patients (41%), the most frequent was Hirschsprung disease was reported in six cases.<sup>40, 44, 59</sup> Neurological diseases were reported in eight cases (18%) and cardiovascular diseases occurred in five patients (11%). Chronic renal failure was reported in six patients (14%).<sup>53, 56, 58, 67, 68</sup> Other conditions found are included in the Tables 1–5.

### Side effects

The side effects mainly included metabolic disturbances, particularly hyperphosphataemia, hypocalcaemia,

hypernatraemia, hypokalaemia and metabolic acidosis. Tetany resulting from the hypocalcaemia-induced was reported in 17 cases (34%),<sup>37, 39, 40, 43, 47, 48, 52–56, 58, 62, 65, 69</sup> being the most frequent complication (the remaining complications can be consulted in Tables 1–5).

### Age

The results in all patients were stratified by age, forming groups of patients under 18 years of age and adults (over 18 years of age). In turn, separate analyses were made in each group of patients <2 years old (in those under 18 years) and patients older than 65 years (in the adult group).

#### *Paediatric age (0–18 years)*

A total of 29 case reports, representing 66% of all reported cases, were identified. Patients aged 2 years or less (15 cases, 34% of all cases analysed)<sup>32–44</sup> and patients older than 2 years (14 cases, 32%)<sup>38, 40, 45–56</sup> were separately analysed. Within the latter group, it should be noted that almost all patients had ages ranging from 1 to 5 years, we only found one case of a child older than 5 years.

#### *Children under 2 years of age*

Nine boys (60%) and six girls (40%) were found under the age of 2. As regards associated comorbidity, an underlying disease was found in 11 cases (73%). Indications for prescription included constipation in six cases (40%)<sup>33, 37, 40</sup> and colon preparation for surgery in another three patients (20%).<sup>32, 36, 39</sup> All cases seen in infants under 2 years of age showed metabolic disturbances. Finally, a case was identified in a newborn that had bone mineralization disturbances probable because of repeated enema used by his anorexic mother during pregnancy<sup>34</sup> (Table 1).

#### *Children aged 2–18 years*

Seven males (50%) and seven females (50%) were found between the ages of 2–18. Twelve of these patients (86%) had associated comorbidity (Table 2). The most common group of conditions were gastrointestinal motility disorders, found in six patients (43%).<sup>38, 40, 47, 49, 50, 54</sup>

Table 1. Adverse effects reported in patients under 2 year of age

Author/year	Sex	Age	Associated condition	Indication	No. of units administered	Disturbances induced	Death
Everman <i>et al.</i> (2003) <sup>32</sup>	Male	1 year	No	Preparation for surgery	1	Water-electrolyte disturbances, acute respiratory failure	No
Ismail <i>et al.</i> (2000) <sup>41</sup>	Male	1.5 years	Asthma, epilepsy	Constipation	1	Water-electrolyte disturbances	Yes
Walton <i>et al.</i> (2000) <sup>33</sup>	Male	6 weeks	Premature birth	Constipation	1	Water-electrolyte disturbances, acute renal failure	Yes
Craig <i>et al.</i> (1994) <sup>42</sup>	Female	2 years	VATER syndrome	Constipation	2	Water-electrolyte disturbances	No
Rimersberger <i>et al.</i> (1992) <sup>34</sup>	Female	Newborn	No	-	Multiple, administered to mother during pregnancy	Bone mineralization disturbances	No
McCabe <i>et al.</i> (1991) <sup>43</sup>	Female	2 years	Cat cry syndrome, heart failure	-	1 (90 mL)	Water-electrolyte disturbances, tetany	No
Wason <i>et al.</i> (1989) <sup>35</sup>	Female	5 months	No	-	1 (adult)	Water-electrolyte disturbances	No
Martin <i>et al.</i> (1987) <sup>36</sup>	Male	11 months	Intestinal reconstruction	Preparation for surgery	4 (adult)	Water-electrolyte disturbances	Yes
Reedy <i>et al.</i> (1983) <sup>37</sup>	Male	1 year	Muscle dystrophy	Constipation	1	Water-electrolyte disturbances, tetany, fever	No
Gómez <i>et al.</i> (1981) <sup>44</sup>	Male	1.3 years	Hirschprung	Constipation	2	Water-electrolyte disturbances	No
Davis <i>et al.</i> (1977) <sup>38</sup>	Female	4 months	Constipation	Constipation	1	Water-electrolyte disturbances	No
Honig <i>et al.</i> (1975) <sup>39</sup>	Male	5 months	Intestinal reconstruction due to imperforate anus	Preparation for surgery	60-mL fleet enema	Water-electrolyte disturbances, tetany, fever	No
Moseley <i>et al.</i> (1968) <sup>40</sup>	Male	8 months	Hirschprung	-	2 every 12 h	Water-electrolyte disturbances, tetany	No
Moseley <i>et al.</i> (1968) <sup>40</sup>	Male	7 months	Hirschprung	-	1/2 adult enema	Water-electrolyte disturbances, tetany	No
Moseley <i>et al.</i> (1968) <sup>40</sup>	Female	2 years	Hirschprung	-	1/2 adult enema	Water-electrolyte disturbances, tetany, QT prolongation	No

Table 2. Adverse effects reported in patients older than 2 years and younger than 18 years

Author/year	Sex	Age (years)	Associated condition	Indication	No. of units administered	Disturbances induced	Death
Butani <i>et al.</i> (2005) <sup>56</sup>	Male	11	Neurogenic bladder, end-stage renal failure	Constipation	2 paediatric enemas	Water-electrolyte disturbances, tetany, QT prolongation	No
Marrafa <i>et al.</i> (2004) <sup>55</sup>	Female	4	Spinal muscular atrophy	Constipation	2 adults enemas within 4 h	Water-electrolyte disturbances, tetany, QT prolongation	No
Melvin <i>et al.</i> (2002) <sup>45</sup>	Male	3	No	Constipation	-	Water-electrolyte disturbances, intravascular haemolysis	No
Ballesteros <i>et al.</i> (2001) <sup>46</sup>	Male	3	Lymphoma, liver transplantation	Constipation	1 (80 mL)	Water-electrolyte disturbances	Yes
Helikson <i>et al.</i> (1997) <sup>47</sup>	Female	3	Anorectal malformation	Constipation	3 (adult)	Water-electrolyte disturbances, tetany	No
Franch <i>et al.</i> (1995) <sup>48</sup>	Female	4	No	Constipation	1 (250 mL)	Water-electrolyte disturbances, tetany	No
Hunter <i>et al.</i> (1993) <sup>49</sup>	Female	4	Constipation	Constipation	2.5 enemas 3 times weekly	Water-electrolyte disturbances, abdominal distention	No
Edmonson <i>et al.</i> (1990) <sup>50</sup>	Male	4	Constipation	Constipation	3 (adult)	Water-electrolyte disturbances, QT prolongation	No
Forman <i>et al.</i> (1979) <sup>51</sup>	Female	3	Gaucher	-	2	Water-electrolyte disturbances	No
Sotos <i>et al.</i> (1977) <sup>52</sup>	Female	3	Myelomeningocele	Constipation	2	Water-electrolyte disturbances, tetany	No
Davis <i>et al.</i> (1977) <sup>38</sup>	Male	3	Constipation	-	1	Neurological disturbances	No
Oxnard <i>et al.</i> (1974) <sup>53</sup>	Male	5	Chronic renal failure, congenital urinary obstruction	-	1 (adult)	Water-electrolyte disturbances, tetany, QT prolongation	No
Swerdlow <i>et al.</i> (1974) <sup>54</sup>	Male	3	Pyloric stenosis	Constipation	1 (undiluted Fosfosoda)	Water-electrolyte disturbances, tetany	No
Moseley <i>et al.</i> (1968) <sup>40</sup>	Female	3	Hirschsprung	-	-	Water-electrolyte disturbances, tetany, QT prolongation	No

**Table 3.** Adverse effects reported in patients aged 18–65 years

Author/year	Sex	Age (years)	Associated condition	Indication	No. of units administered	Disturbances induced	Death
Eckstein <i>et al.</i> (2006) <sup>60</sup>	Female	64	Kidney transplantation, hyperparathyroidism, gastrectomy	Constipation	3 (in several days)	Extensive calcifications, liver enzymes elevation, multiorgan failure and shock	Yes
Pitcher <i>et al.</i> (1997) <sup>57</sup>	Male	64	Rectal neoplasm	Preparation for colonoscopy	–	Water-electrolyte disturbances, shock, multiorgan failure	Yes
Haskell <i>et al.</i> (1985) <sup>58</sup>	Male	58	Polycystic renal disease, chronic renal failure	Constipation	2	Water-electrolyte disturbances, tetany	No
Young <i>et al.</i> (1968) <sup>59</sup>	Male	21	Hirschsprung	Constipation	4 in 48 h	Water-electrolyte disturbances	No

The indication for enema was constipation in 10 cases (71%),<sup>45–50, 52, 54–56</sup> while no data on indication was found in the remaining four patients (35%).<sup>38, 40, 51, 53</sup> All patients experienced the previously reported water and electrolyte disturbances. Other conditions included tetany in seven cases (50%)<sup>40, 47, 52–56</sup> and QT interval prolongation in five cases (36%).<sup>40, 50, 53, 55, 56</sup> One death (7%) was identified in this group, in a male with a significant comorbidity (gastrointestinal lymphoma and liver transplantation).<sup>46</sup>

#### Adults aged 18–65 years.

In the adult group, a total of 15 cases with adverse effects were found. Four of these occurred in patients under 65 years of age,<sup>57–60</sup> and 11 in patients over 65 years of age.<sup>61–70</sup> The mean age in patients aged 18–65 years who experienced adverse effects was 52 years. They all had comorbidity of a different severity. Water and electrolyte disturbances occurred in all cases. We found two deaths in this group<sup>57, 60</sup> (Table 3).

#### Adults over 65 years of age

Finally, 11 clinical cases, eight females and three males with a mean age of 81 years (range: 70–96 years), were found among patients older than 65 years. They all had comorbid conditions. The most common associated conditions were heart diseases, reported in six patients (55%).<sup>61–64, 67, 68</sup> Indications included constipation in nine cases (73%)<sup>61–63, 65–68, 70</sup> and preparation for colonoscopy or barium enema in two patients (27%).<sup>64, 69</sup> As regards the dosage given, seven patients (64%) received three or more units,<sup>61–63, 65, 66, 69, 70</sup> and a maximum of eight doses were received by a single patient.<sup>62</sup> All patients experienced water and electrolyte disturbances (Table 4). Six patients over 65 years of age died (55%).<sup>61–63, 67, 70</sup> Five of these patients (45%) had been given three or more doses,<sup>61–63, 70</sup> and the remaining patient had significant associated comorbidity (acute pulmonary oedema, heart failure and chronic renal failure).<sup>67</sup>

#### Mortality

Among all aforementioned studies, a total of 12 deaths (27%) were found, six in males and six in females.<sup>33, 36, 41, 46, 57, 60–63, 67, 70</sup> Eleven of such deaths

Table 4. Adverse effects reported in patients over 65 years of age

Author/year	Sex	Age (years)	Associated condition	Indication	No. of units administered	Disturbances induced	Death
Farah <i>et al.</i> (2005) <sup>70</sup>	Male	70	Spondyloarthropathy	Constipation	4 enemas within 12 h (133 ml each)	Water-electrolyte disturbances, cardiac arrest	Yes
Tan <i>et al.</i> (2002) <sup>61</sup>	Female	73	Heart failure	Constipation	3	Water-electrolyte disturbances	Yes
Tan <i>et al.</i> (2002) <sup>61</sup>	Female	82	-	Constipation	3	Water-electrolyte disturbances	Yes
Martinez Velasco <i>et al.</i> (1998) <sup>62</sup>	Female	86	Atrial fibrillation	Constipation	8	Water-electrolyte disturbances, tetany	Yes
Knobel <i>et al.</i> (1996) <sup>63</sup>	Female	87	Ischaemic heart disease, high blood pressure, megacolon	Constipation	4 in 48 h	Water-electrolyte disturbances, coma, and respiratory failure	Yes
Sutters <i>et al.</i> (1996) <sup>64</sup>	Male	71	Chronic obstructive pulmonary disease, high blood pressure, supraventricular tachycardia	Preparation for colonoscopy	2	Water-electrolyte disturbances	No
Korzets <i>et al.</i> (1992) <sup>65</sup>	Female	77	Urinary incontinence	Constipation	6 in 12 h	Water-electrolyte disturbances, confusion, QT prolongation, and tetany	No
Aradhya <i>et al.</i> (1991) <sup>66</sup>	Female	96	Dementia, gastrostomy	Constipation	2	Water-electrolyte disturbances, lethargy	No
Spinrad <i>et al.</i> (1989) <sup>67</sup>	Female	91	Heart failure, acute pulmonary oedema, chronic renal failure	Constipation	1	Water-electrolyte disturbances	Yes
Biberstein <i>et al.</i> (1985) <sup>68</sup>	Male	81	Chronic renal failure, atrial fibrillation, atherosclerosis	Constipation	1	Water-electrolyte disturbances, QT prolongation	No
Rohack <i>et al.</i> (1985) <sup>69</sup>	Female	77	Diverticulitis	Barium enema preparation	6 in 6 h	Water-electrolyte disturbances, coma, tetany, and fever	No

Table 5. Dead patients

Author/year	Sex	Age	Associated condition	Indication	No of units administered	Cause of death
Eckstein <i>et al.</i> (2006) <sup>60</sup>	Female	64 years	Kidney transplantation, hyperparathyroidism, gastrectomy	Constipation	3 (in several days)	Extensive calcifications, liver enzymes elevation, multiorgan failure and shock
Farah <i>et al.</i> (2005) <sup>70</sup>	Male	70 years	Spondylortopathy	Constipation	4 enemas within 12 h (133 ml each)	Water-electrolyte disturbances, cardiac arrest
Tan <i>et al.</i> (2002) <sup>61</sup>	Female	73 years	Heart failure	Constipation	3	Pneumonia. Water-electrolyte disturbances
Tan <i>et al.</i> (2002) <sup>61</sup>	Female	82 years	No	Constipation	3	Water-electrolyte disturbances
Ballesteros Garcia <i>et al.</i> (2001) <sup>46</sup>	Male	3 years	Lymphoma, liver transplantation	Constipation	1 (80 mL)	Water-electrolyte disturbances
Ismail <i>et al.</i> (2000) <sup>41</sup>	Male	17 months	Asthma, epilepsy	Constipation	1	Water-electrolyte disturbances
Walton <i>et al.</i> (2000) <sup>33</sup>	Male	6 weeks	Premature birth	Constipation	1	Water-electrolyte disturbances, acute renal failure
Martinez Velasco <i>et al.</i> (1998) <sup>62</sup>	Female	86 years	Atrial fibrillation	Constipation	8	Water-electrolyte disturbances
Pitcher <i>et al.</i> (1997) <sup>57</sup>	Male	64 years	Rectal neoplasm	Preparation for colonoscopy	-	Water-electrolyte disturbances
Knobel <i>et al.</i> (1996) <sup>63</sup>	Female	87 years	High blood pressure, ischaemic heart disease	Constipation	4	Water-electrolyte disturbances
Spinard <i>et al.</i> (1989) <sup>67</sup>	Female	91 years	Heart failure, acute pulmonary oedema, chronic renal failure	Constipation	1	Water-electrolyte disturbances
Martin <i>et al.</i> (1987) <sup>36</sup>	Male	11 months	Imperforate anus, colostomy	Preparation for surgery	4	Water-electrolyte disturbances

(92%) were due to water and electrolyte disturbances secondary to administration of sodium phosphate enemas. Age of dead patients ranged from 11 months<sup>36</sup> to 91 years.<sup>67</sup> Virtually, all dead patients were at the extreme ages of life. Four deaths (30%)<sup>33, 36, 41, 46</sup> occurred among patients under 18 years of age. In the group of patients over 18 years of age, a total of six deaths (55%)<sup>57, 60–63, 67, 70</sup> occurred in patients aged 64–91 years. Among adult patients aged 18–65 years, death was only reported in two clinical cases (Table 5).

The subgroup of adults who died included six women (75%)<sup>60–62, 67</sup> and two men.<sup>57, 70</sup> All patients under 18 years of age who died were males.<sup>33, 36, 41, 46</sup>

All dead patients except 1 (92%)<sup>61</sup> had associated comorbidity in all age groups studied. (Table 5).

## DISCUSSION

Sodium phosphate enemas are products widely used in both in-patient and out-patient settings. The most common indication is for symptomatic treatment of constipation, and to a lesser extent in preparation for colonoscopy or surgery. There are no accurate data about worldwide prescription of these products. According to the manufacturer (Casen/Fleet), more than 5 00 000 000 U have been sold up to now, which can give us an approximate idea of the widespread use of these products.

No randomized clinical trials, meta-analysis or systematic reviews exist in the literature to answer the question of safeness or adverse effects of these products. The side effects are minimal, and literature reports only refer to the most severe cases, such as water and electrolyte disturbances that may even be fatal in some cases. There are various randomized clinical trials comparing the tolerability and efficacy for colon cleaning of several preparation methods for endoscopic procedures. Such trials assessed the side effects of phosphate enemas that were considered to be mild and with no clinical impact. In a US study<sup>3</sup> conducted on 157 patients, nausea (6–18%), vomiting (0–7%) and abdominal pain (8–9%) were reported, while abdominal distention occurred in 90% of subjects receiving one enema and in 98% of patients when two enemas were administered. Atkin *et al.*<sup>2</sup> found similar results in an analysis of 721 patients receiving a sodium phosphate enema.

It should not be forgotten that sodium phosphate enemas are widely used, and our review only found a

minimum number of patients with side effects. Specifically, the review conducted found 46 reports of side effects of different severity, which would represent a minimal proportion of side effects if we take into account the widespread use of enemas. Nevertheless, these data should be analysed with caution because of the possible publication bias incurred, as only a minority of side effects may be reported, and they are probably the most severe.

As regards age distribution of patients reported as experiencing side effects, it should be noted that most of them were in the extreme age groups (older than 65 years and younger than 5 years, 25% and 64%, respectively). Only five cases were reported in patients aged from 5 to 65 years. It should therefore be inferred that extreme ages are associated with a greater frequency of side effects.

Comorbid conditions were noted in 86% of cases, particularly including neurological, gastrointestinal and renal disorders. Such associated conditions could be related to the increased phosphate absorption shown in some clinical trials. Thus, in the Schumann *et al.* study<sup>28</sup>, high serum phosphorus levels were shown in patients with a longer enema retention time. There have been reports of several experimental studies in animals showing phosphorus absorption in the colonic mucosa,<sup>29, 30</sup> that is dependent on luminal phosphorus levels. By contrast, other studies analysing water and electrolyte disturbances in patients who were prepared for colonoscopy using sodium phosphate enemas only showed a mild increase in serum phosphate levels that did not reach pathological ranges.<sup>26, 28</sup> It could be hypothesized that the existence of increased blood phosphate levels in patients with gastrointestinal disorders could be due to an increased contact between enema contents and the intestinal wall, which would promote phosphorus and sodium absorption.

The actual dosage administered to cases reported in the literature is difficult to assess, as neither the dose nor the composition of enemas are adequately reported in most publications. It should also be noted that formulations differ depending on the country.<sup>75</sup> Moreover, some publications report out-patient administration of adult enemas to paediatric patients. Overdosage was reported in eight clinical cases, and up to eight enemas were administered to a 86-year-old patient.<sup>62</sup> In this respect, it should be noted that most dead patients had been administered two or more enemas. An influence of the dose



received on adverse effects and their severity is therefore likely.

The side effects reported are related to water and electrolyte disturbances resulting from hyperphosphatemia, hypocalcaemia, hypernatraemia, and metabolic acidosis, because of the absorptive effect of enema components and to their inadequate elimination in some cases, such as patients with chronic renal failure. Therefore, an increased caution is required when enemas are administered to patients with this condition.

It should be noted that virtually all deaths reported in the literature occurred in people with extreme ages and a significant comorbidity. Deaths were caused by water and electrolyte disturbances, by an episode of pneumonia and by an extensive calcification with subsequent multiorgan failure in a patient with renal transplantation and hyperparathyroidism. To sum up, age older than 65 years and under 5 years could be suggested as a potential risk factor for mortality, which could be related to an increase in associated conditions.

To summarize, it may be concluded that water and electrolyte disturbances are the reason for the main side effects occurring in patients administered sodium phosphate enemas. The main risk factors include chronic renal failure, diseases altering intestinal motility (neurological, morphological, etc.), and extreme ages of life. Adequate prescription is required in patients with such conditions, as use of sodium phosphate enemas are not risk-free, although the incidence of side effects is nevertheless very low. By contrast, administration of sodium phosphate enemas does not involve a serious risk for health in patients without such risk factors, who represent the majority of cases.

## ACKNOWLEDGMENTS

*Declaration of personal interests:* None. *Declaration of funding interests:* this review was funded in part by the Instituto de Salud Carlos III (grant numbers C03/02 and PI050109)

## REFERENCES

- Harrington L, Schuh S. Complications of Fleet enema administration and suggested guidelines for use in the pediatric emergency department. *Pediatr Emerg Care* 1997; 13: 225–6.
- Atkin WS, Hart A, Edwards R, *et al.* Single blind, randomised trial of efficacy and acceptability of oral picolax versus self administered phosphate enema in bowel preparation for flexible sigmoidoscopy screening. *BMJ* 2000; 320: 1504–8; discussion 1509.
- Osgard E, Jackson JL, Strong J. A randomized trial comparing three methods of bowel preparation for flexible sigmoidoscopy. *Am J Gastroenterol* 1998; 93: 1126–30.
- Bowers B. Evaluating the evidence for administering phosphate enemas. *Br J Nurs* 2006; 15: 378–81.
- Laxatives for bowel clearing before investigations. *Drug Ther Bull* 2002; 40: 86–8.
- Herman M, Shaw M, Loewen B. Comparison of three forms of bowel preparations for screening flexible sigmoidoscopy. *Gastroenterol Nurs* 2001; 24: 178–81.
- Toledo TK, DiPalma JA. Review article: colon cleansing preparation for gastrointestinal procedures. *Aliment Pharmacol Ther* 2001; 15: 605–11.
- Fincher RK, Osgard EM, Jackson JL, *et al.* A comparison of bowel preparations for flexible sigmoidoscopy: oral magnesium citrate combined with oral bisacodyl, one hypertonic phosphate enema, or two hypertonic phosphate enemas. *Am J Gastroenterol* 1999; 94: 2122–7.
- Manoucheri M, Nakamura DY, Lukman RL. Bowel preparation for flexible sigmoidoscopy: which method yields the best results? *J Fam Pract* 1999; 48: 272–4.
- Preston KL, Peluso FE, Goldner F. Optimal bowel preparation for flexible sigmoidoscopy—are two enemas better than one? *Gastrointest Endosc* 1994; 40: 474–6.
- Barrish JO, Gilger MA. Colon cleanout preparations in children and adolescents. *Gastroenterol Nurs* 1993; 16: 106–9.
- Krevsky B, Niewiarowski T, League R, *et al.* Flexible sigmoidoscopy screening in an industrial setting. *Am J Gastroenterol* 1992; 87: 1759–62.
- Sugimura F, Ryoh H, Watanabe T, *et al.* Comparative studies on the usefulness of phosphate versus glycerin enema in preparation for colon examinations. *Gastroenterol Jpn* 1990; 25: 437–50.
- Richter KP, Cleveland MB. Comparison of an orally administered gastrointestinal lavage solution with traditional enema administration as preparation for colonoscopy in dogs. *J Am Vet Med Assoc* 1989; 195: 1727–31.
- Mulder CJ, Tytgat GN, Wiltink EH, *et al.* Comparison of 5-aminosalicylic acid (3 g) and prednisolone phosphate sodium enemas (30 mg) in the treatment of distal ulcerative colitis. A prospective, randomized, double-blind trial. *Scand J Gastroenterol* 1988; 23: 1005–8.
- Mulder CJ, Endert E, van der Heide H, *et al.* Comparison of beclomethasone dipropionate (2 and 3 mg) and prednisolone sodium phosphate enemas (30 mg) in the treatment of ulcerative proctitis. An adrenocortical approach. *Neth J Med* 1989; 35: 18–24.
- van der Heide H, van den Brandt-Gradel V, Tytgat GN, *et al.* Comparison of beclomethasone dipropionate and prednisolone 21-phosphate enemas in

- the treatment of ulcerative proctitis. *J Clin Gastroenterol* 1988; 10: 169–72.
- 18 Jorgensen LS, Center SA, Randolph JF, et al. Electrolyte abnormalities induced by hypertonic phosphate enemas in two cats. *J Am Vet Med Assoc* 1985; 187: 1367–8.
  - 19 Somerville KW, Langman MJ, Kane SP, et al. Effect of treatment on symptoms and quality of life in patients with ulcerative colitis: comparative trial of hydrocortisone acetate foam and prednisolone 21-phosphate enemas. *Br Med J (Clin Res Ed)* 1985; 291: 866.
  - 20 Atkins CE, Tyler R, Greenlee P. Clinical, biochemical, acid-base, and electrolyte abnormalities in cats after hypertonic sodium phosphate enema administration. *Am J Vet Res* 1985; 46: 980–8.
  - 21 McCallum RW, Meyer CT, Marignani P, et al. Flexible sigmoidoscopy: diagnostic yield in 1015 patients. *Am J Gastroenterol* 1984; 79: 433–7.
  - 22 Lambert R, Olive C, Melange M, et al. Flexible rectosigmoidoscopy in the detection of tumoral colonic lesions. *Endoscopy* 1978; 10: 284–8.
  - 23 Powell-Tuck J, Lennard-Jones JE, May CS, et al. Plasma prednisolone levels after administration of prednisolone-21-phosphate as a retention enema in colitis. *Br Med J* 1976; 1: 193–5.
  - 24 Pinfield A, Stringer MD. Randomised trial of two pharmacological methods of bowel preparation for day case colonoscopy. *Arch Dis Child* 1999; 80: 181–3.
  - 25 Dahshan A, Lin CH, Peters J, et al. A randomized, prospective study to evaluate the efficacy and acceptance of three bowel preparations for colonoscopy in children. *Am J Gastroenterol* 1999; 94: 3497–501.
  - 26 Cohan CF, Kadakia SC, Kadakia AS. Serum electrolyte, mineral, and blood pH changes after phosphate enema, water enema, and electrolyte lavage solution enema for flexible sigmoidoscopy. *Gastrointest Endosc* 1992; 38: 575–8.
  - 27 Platell C, Barwood N, Makin G. Randomized clinical trial of bowel preparation with a single phosphate enema or polyethylene glycol before elective colorectal surgery. *Br J Surg* 2006; 93: 427–33.
  - 28 Schuchmann GD, Barcia PJ. Phosphate absorption from fleet enemas in adults. *Curr Surg* 1989; 46: 120–2.
  - 29 Hu MS, Kayne LH, Jamgotchian N, et al. Paracellular phosphate absorption in rat colon: a mechanism for enema-induced hyperphosphatemia. *Miner Electrolyte Metab* 1997; 23: 7–12.
  - 30 Lochbuhler H, Sachs J, Raute-Kreinsen U. The pharmacological effect of sodium phosphate after absorption from the peritoneal cavity. *Eur J Pediatr Surg* 1995; 5: 84–7.
  - 31 Gutierrez E. Purgative with high sodium phosphate contents: efficacious but not so safe. *Med Clin (Barc)* 2006; 126: 173–4.
  - 32 Everman DB, Nitu ME, Jacobs BR. Respiratory failure requiring extracorporeal membrane oxygenation after sodium phosphate enema intoxication. *Eur J Pediatr* 2003; 162: 517–9.
  - 33 Walton DM, Thomas DC, Aly HZ, et al. Morbid hypocalcemia associated with phosphate enema in a six-week-old infant. *Pediatrics* 2000; 106: E37.
  - 34 Rimensberger P, Schubiger G, Willi U. Connatal rickets following repeated administration of phosphate enemas in pregnancy: a case report. *Eur J Pediatr* 1992; 151: 54–6.
  - 35 Wason S, Tiller T, Cunha C. Severe hyperphosphatemia, hypocalcemia, acidosis, and shock in a 5-month-old child following the administration of an adult Fleet enema. *Ann Emerg Med* 1989; 18: 696–700.
  - 36 Martin RR, Lisehora GR, Braxton M Jr, et al. Fatal poisoning from sodium phosphate enema. Case report and experimental study. *JAMA* 1987; 257: 2190–2.
  - 37 Reedy JC, Zwiren GT. Enema-induced hypocalcemia and hyperphosphatemia leading to cardiac arrest during induction of anesthesia in an outpatient surgery center. *Anesthesiology* 1983; 59: 578–9.
  - 38 Davis RF, Eichner JM, Bleyer WA, et al. Hypocalcemia, hyperphosphatemia, and dehydration following a single hypertonic phosphate enema. *J Pediatr* 1977; 90: 484–5.
  - 39 Honig PJ, Holtzapfel PG. Hypocalcemic tetany following hypertonic phosphate enemas. *Clin Pediatr (Phila)* 1975; 14: 678–9.
  - 40 Moseley PK, Segar WE. Fluid and serum electrolyte disturbances as a complication of enemas in Hirschsprung's disease. *Am J Dis Child* 1968; 115: 714–8.
  - 41 Ismail EA, Al-Mutairi G, Al-Anzy H. A fatal small dose of phosphate enema in a young child with no renal or gastrointestinal abnormality. *J Pediatr Gastroenterol Nutr* 2000; 30: 220–1.
  - 42 Craig JC, Hodson EM, Martin HC. Phosphate enema poisoning in children. *Med J Aust* 1994; 160: 347–51.
  - 43 McCabe M, Sibert JR, Routledge PA. Phosphate enemas in childhood: cause for concern. *BMJ* 1991; 302: 1074.
  - 44 Gomez Rivas B, Labay Matias M, Reynes Muntaner J, et al. Hypocalcemic tetany caused by a phosphate enema. *An Esp Pediatr* 1981; 14: 143–4.
  - 45 Melvin JD, Watts RG. Severe hypophosphatemia: a rare cause of intravascular hemolysis. *Am J Hematol* 2002; 69: 223–4.
  - 46 Ballesteros Garcia M, Sanchez Diaz JI, Mar Molinero F. Poisoning after the use of sodium phosphate enema. *An Esp Pediatr* 2001; 55: 92–3.
  - 47 Helikson MA, Parham WA, Tobias JD. Hypocalcemia and hyperphosphatemia after phosphate enema use in a child. *J Pediatr Surg* 1997; 32: 1244–6.
  - 48 Franch F, Verd F, Hernandez P, et al. Hypocalcemic tetany following the administration of phosphate enema. *Rev Esp Anestesiol Reanim* 1995; 42: 35–6.
  - 49 Hunter MF, Ashton MR, Griffiths DM, et al. Hyperphosphatemia after enemas in childhood: prevention and treatment. *Arch Dis Child* 1993; 68: 233–4.
  - 50 Edmondson S, Almqvist TD. Iatrogenic hypocalcemic tetany. *Ann Emerg Med* 1990; 19: 938–40.
  - 51 Forman J, Baluarte HJ, Gruskin AB. Hypokalemia after hypertonic phosphate enemas. *J Pediatr* 1979; 94: 149–51.
  - 52 Sotos JF, Cutler EA, Finkel MA, et al. Hypocalcemic coma following two pediatric phosphate enemas. *Pediatrics* 1977; 60: 305–7.
  - 53 Oxnard SC, O'Bell J, Grupe WE. Severe tetany in an azotemic child related to a sodium phosphate enema. *Pediatrics* 1974; 53: 105–6.
  - 54 Swerdlow DB, Labow S, D'Anna J. Tetany and enemas: report of a case. *Dis Colon Rectum* 1974; 17: 786–7.
  - 55 Marraffa JM, Hui A, Stork CM. Severe hyperphosphatemia and hypocalcemia following the rectal administration of a phosphate-containing Fleet pediatric enema. *Pediatr Emerg Care* 2004; 20: 453–6.
  - 56 Butani L. Life-threatening hyperphosphatemia and hypocalcemia from inappropriate use of Fleet enemas. *Clin Pediatr (Phila)* 2005; 44: 93.
  - 57 Pitcher DE, Ford RS, Nelson MT, et al. Fatal hypocalcemic, hyperphosphatemic, metabolic acidosis following sequential sodium phosphate-based enema admin-

- istration. *Gastrointest Endosc* 1997; **46**: 266–8.
- 58 Haskell LP. Hypocalcaemic tetany induced by hypertonic-phosphate enema. *Lancet* 1985; **2**: 1433.
- 59 Young JF, Brooke BN. Enema shock in Hirschsprung's disease. *Dis Colon Rectum* 1968; **11**: 391–5.
- 60 Eckstein J, Savic S, Eugster T, *et al.* Extensive calcifications induced by hyperphosphataemia caused by phosphate-based enema in a patient after kidney transplantation. *Nephrol Dial Transplant* 2006; **21**: 2013–6.
- 61 Tan HL, Liew QY, Loo S, *et al.* Severe hyperphosphataemia and associated electrolyte and metabolic derangement following the administration of sodium phosphate for bowel preparation. *Anaesthesia* 2002; **57**: 478–83.
- 62 Martinez Velasco MC, Ahmad al Ghool M, Sos Ortigosa F, *et al.* Acute hyperphosphatemia induced by enemas. *Med Clin (Barc)* 1998; **110**: 805.
- 63 Knobel B, Petchenko P. Hyperphosphatemic hypocalcemic coma caused by hypertonic sodium phosphate (fleet) enema intoxication. *J Clin Gastroenterol* 1996; **23**: 217–9.
- 64 Sutters M, Gaboury CL, Bennett WM. Severe hyperphosphatemia and hypocalcemia: a dilemma in patient management. *J Am Soc Nephrol* 1996; **7**: 2056–61.
- 65 Korzets A, Dicker D, Chaimoff C, *et al.* Life-threatening hyperphosphatemia and hypocalcemic tetany following the use of fleet enemas. *J Am Geriatr Soc* 1992; **40**: 620–1.
- 66 Aradhye S, Brensilver JM. Sodium phosphate-induced hypernatremia in an elderly patient: a complex pathophysiologic state. *Am J Kidney Dis* 1991; **18**: 609–11.
- 67 Spinrad S, Sztern M, Grosskopf Y, *et al.* Treating constipation with phosphate enema: an unnecessary risk. *Isr J Med Sci* 1989; **25**: 237–8.
- 68 Biberstein M, Parker BA. Enema-induced hyperphosphatemia. *Am J Med* 1985; **79**: 645–6.
- 69 Rohack JJ, Mehta BR, Subramanyam K. Hyperphosphatemia and hypocalcemic coma associated with phosphate enema. *South Med J* 1985; **78**: 1241–2.
- 70 Farah R. Fatal acute sodium phosphate enemas intoxication. *Acta Gastroenterol Belg* 2005; **68**: 392–3.
- 71 Smith I, Carr N, Corrado OJ, *et al.* Rectal necrosis after a phosphate enema. *Age Ageing* 1987; **16**: 328–30.
- 72 Sweeney JL, Hewett P, Riddell P, *et al.* Rectal gangrene: a complication of phosphate enema. *Med J Aust* 1986; **144**: 374–5.
- 73 Pietsch JB, Shizgal HM, Meakins JL. Injury by hypertonic phosphate enema. *Can Med Assoc J* 1977; **116**: 1169–70.
- 74 Bell AM. Colonic perforation with a phosphate enema. *J R Soc Med* 1990; **83**: 54–5.
- 75 Post SS. Hyperphosphatemic hypocalcemic coma caused by hypertonic sodium phosphate (fleet) enema intoxication. *J Clin Gastroenterol* 1997; **24**: 192.

# Differences in characteristics of nocturnal enuresis between children and adolescents: a critical appraisal from a large epidemiological study

CHUNG K. YEUNG, BIJI SREEDHAR, JENNIFER D.Y. SIHOE, FRANCES K.Y. SIT and JOSEPH LAU\*

Department of Surgery, Division of Paediatric Surgery and Paediatric Urology, and \*Centre for Clinical Trials and Epidemiological Research, Chinese University of Hong Kong, Prince of Wales Hospital, Hong Kong

Accepted for publication 15 November 2005

## OBJECTIVE

To evaluate any differences in the characteristics of primary nocturnal enuresis (PNE) between younger enuretic children and adolescents.

## SUBJECTS AND METHODS

In all, 21 000 questionnaires designed to determine the presence or absence of bed-wetting, diurnal incontinence, frequency of wetting, systemic illness, and family history, were sent to children aged 5–19 years from 67 kindergartens, primary schools and secondary schools randomly selected by a computer from different areas in Hong Kong. In addition, questions were asked to evaluate when and how the parents became aware that bed-wetting is a significant medical problem deserving attention in children after the age of 5 years.

## RESULTS

Of the 21 000 questionnaires distributed, 16 512 (78.6%) were completed. Among the

respondents, 512 children (302 boys, 210 girls) had PNE; of these, 106 (20.7%) also had daytime incontinence. There was a marked reduction in the overall prevalence of PNE with advancing age. At 5 years old, 16.1% of children had PNE (20.7% boys, 10.8% girls; at age 9 and 19 years, 3.14% and 2.2% of children had PNE, respectively. However, this reduction was significantly more apparent among those with mild enuretic symptoms (wet <3 nights/week) than in those with more frequent bed-wetting. Furthermore, younger enuretic children behaved very differently from adolescents and older patients. As age increased there was a significant tendency towards more severe enuretic symptoms. At age 5 years, 14.3% of enuretic children wet 7 nights/week, compared with 48.3% at age 19 years ( $P < 0.001$ ). In addition, significantly more adolescent boys aged >10 years had daytime urinary incontinence than had enuretic children aged  $\leq 10$  years (32% vs 14.6%, respectively,  $P < 0.001$ ). Most (89%) parents only became aware that bed-wetting was a significant medical problem deserving attention through material in the mass media over the past 3–4 years.

## CONCLUSIONS

The present finding suggesting that PNE spontaneously resolves with increasing age probably applies only to those with mild enuretic symptoms. There are significant differences in characteristics between younger enuretic children and older subjects. As age increases there is an increasing proportion of enuretic patients with more severe bed-wetting. Enuretic children aged >10 years and adolescents have significantly more daytime urinary symptoms and incontinence. The previously reported low prevalence of PNE in Hong Kong was probably due to parental indifference to the problem.

## KEYWORDS

primary nocturnal enuresis, adolescents, daytime symptoms

## INTRODUCTION

Primary nocturnal enuresis (PNE), or bed-wetting, is a very common clinical and a significant social problem in childhood and adolescence. From published epidemiological studies it is apparent that although the values vary somewhat among different countries, the overall prevalence of PNE remains relatively constant, irrespective of geographical locations. It was estimated that 20–25% of children by the age of 4 years and 10% of children at 7 years are frequent bed-wetters [1–5]. In general, the prevalence decreases with increasing age.

However, the condition often has a profound psychological and social impact on the affected children and their families, and generates significant anxiety and even conflicts among them.

A previous epidemiological survey for PNE in Hong Kong in 1995 indicated that the prevalence of PNE in local Chinese school children was surprisingly lower than all reported series from elsewhere [6]. Since then, public educational programmes both in Hong Kong and in surrounding Asian countries have been introduced. In addition, mass screening for PNE in Hong Kong

through the local Student Health Service has been implemented. Concurrently, in a tertiary referral centre for PNE, we have seen a rapid increase in severely enuretic children, many of whom have never been regarded by their parents as having a medical problem until very recently. Similarly, results of the subsequent epidemiological studies in other Asian countries indicate a much higher prevalence of PNE than that reported previously in Hong Kong, and were similar to values reported in other western countries. To re-evaluate the actual prevalence of PNE among local children, a repeat epidemiological study

involving a much larger sample was therefore conducted.

**SUBJECTS AND METHODS**

Children aged 5–19 years from 67 kindergartens, primary schools and secondary schools, with a greater emphasis on adolescents and teenage groups, were randomly selected among different areas of Hong Kong. A self-administered questionnaire comprising two sections, identical to that used in our previous study, was designed [6]. The first part of the questionnaire determined the demographic details, e.g. age, sex, and family history of enuresis; the second part included questions about the details of bed-wetting problems, if any. Questions included were the presence of daytime wetting, night-time wetting and frequency of wetting. In addition, questions were set to ask the parents when and how they became aware that bed-wetting was a significant medical problem that deserves attention in children after the age of 5 years.

Based on findings from our previous epidemiological study, and given an expectation that up to 2.5–3% of children and adolescents in Hong Kong are likely to have PNE, 21 000 questionnaires were sent to different schools, to ensure an adequate number of enuretic subjects for further statistical analysis. Sealed envelopes were provided to return the questionnaires and to ensure confidentiality. An information leaflet was attached to the questionnaire informing the parent of the voluntary nature of the study. All unreturned and incomplete responses were considered as nonresponders. Enuresis was defined as having at least one wet night every 3 months, as described previously [6]. To evaluate the factors associated with the severity of enuresis, subjects with enuresis were stratified by age. The frequency of bed-wetting was divided into three severity subgroups as <3, 3–6 or 7 wet nights/week, respectively.

The chi-square exact probability test was used to compare the prevalence in PNE between boys and girls in corresponding age groups, and to compare the severity of PNE and associated day symptoms in children and adolescents. To identify significant differences in the prevalence of PNE in boys and girls between different age groups, the chi-square test for trend was used, with  $P < 0.05$  considered to indicate statistical significance.

FIG. 1. Prevalence of PNE in Hong Kong schoolchildren.




TABLE 1 The prevalence of PNE in Hong Kong school children

Age, years	Male enuretics/normal	Female enuretics/normal	Total enuretics/normal	Mean % (95% CI)
5	48/232	22/203	70/435	16.11 (12.6–19.50)
6	38/212	20/236	58/448	12.95 (9.8–16.10)
7	36/335	23/249	59/584	10.1 (7.7–12.5)
8	19/244	17/342	36/586	6.14 (4.2–8.1)
9	19/445	11/523	30/968	3.14 (2.0–4.2)
10	11/365	15/621	26/986	2.63 (1.6–3.6)
11	16/485	7/573	23/1058	2.17 (1.3–3.1)
12	15/433	5/554	20/987	2.02 (1.1–2.9)
13	10/506	6/520	16/1026	1.55 (0.8–2.3)
14	8/523	7/481	15/1004	1.49 (0.7–2.2)
15	7/621	8/546	15/1167	1.28 (0.6–1.9)
16	22/1014	16/1277	38/2291	1.65 (1.1–2.2)
17	20/803	17/1040	37/1843	2.0 (1.4–2.6)
18	19/768	21/1060	40/1828	2.19 (1.5–2.9)
19	14/469	15/832	29/1301	2.23 (1.4–3.0)
Total	302/7455	210/9057	512/16512	3.10 (2.8–3.4)

**RESULTS**

Of the 21 000 questionnaires distributed, 16 512 (78.6%) were completed, from 7455 (45.1%) boys and 9057 (54.9%) girls (mean age 13.68 years). Among these, 512 children had PNE, giving an overall prevalence of 3.1% (4.0% boys and 2.31% girls); of these 512, 106 (20.7%) also had daytime urinary incontinence. There was a marked reduction in the overall prevalence of PNE with increasing age (Fig. 1). At age 5 years, 16.1% (20.7% boys, 10.8% girls) children had PNE, at

7 years, 10.1% (10.7% and 9.2%, respectively) and at 9 and 19 years, 3.14% and 2.2% had PNE, respectively (Table 1). Table 1 also shows that there was a significant and decreasing trend in the prevalence of PNE in both genders with age (both  $P < 0.001$ ).

Although there was a marked reduction in the overall prevalence of PNE with increasing age, it was significantly more apparent among those with mild enuretic symptoms (<3 wet nights/week) than in those with more frequent bed-wetting (Fig. 2). Overall, 82% of

FIG. 2. Severity of PNE in children.





FIG. 3. Severity of PNE vs age.



the adolescent subjects had >3 wet nights/week vs enuretic children aged 5–10 years (42.3%,  $P < 0.001$ , Table 2). As age increased there was a greater proportion of enuretic subjects with more severe bed-wetting. Daytime urinary incontinence was significantly more prevalent in adolescents than enuretic children (Table 2). Moreover, the prevalence of daytime urinary incontinence was significantly greater in adolescent boys than in boys aged 5–10 years (32% vs 14.6%, respectively,  $P < 0.001$ ). In general, enuretic symptoms in the adolescent subjects were more severe than those in children. At age 5 years, only 14.3% of enuretic children wet 7 nights/week, compared with 48.3% at age 19 years ( $P < 0.001$ ; Fig. 3).

Of the children with PNE, 39% considered that PNE was not a serious problem, while 61% of the children considered it as a serious problem with a significant impact on their daily life. Among these children, 28% became aware of this problem in the last 6 months, 21% had been aware for 1–2 years, 4% for 3–4 years and only 7% for >5 years.

DISCUSSION

Nocturnal enuresis is an old but still prevalent clinical problem in childhood and adolescence. The traditional view is that in most cases bed-wetting is due to a developmental immaturity of voiding control, and most enuretic children will ultimately acquire normal control with age. However, previous studies showed that, although spontaneous resolution can continue throughout childhood and adolescence, enuretic problems may persist in 1.5–3% of the adult population [7–10]. Previous studies also show that the frequency and severity of wetting episodes progressively increases with age; those with severe symptoms are much more likely to have persistent problems into adult life [11,12].

A previous epidemiological study for PNE in Hong Kong revealed a low prevalence in local Chinese schoolchildren; at 5 years old, only 10.4% of boys and 6.6% of girls had PNE. The prevalence decreased rapidly with increasing age and by 7 and 10 years old, only 4.9% and 1.2% of boys and 0.5% and no girls had enuresis, respectively [6]. These values were significantly lower than all previously published series from other countries [1–5,13–18].

TABLE 2 The severity of NE in younger children and adolescents, and the incidence of daytime urinary incontinence in enuretic younger children and adolescents, as n (%)

Condition	Children (5–10 years)	Adolescents (>10 years)	Total	P
<b>Severity of PNE</b>				
<3 wet nights/week	161/279 (57.7)	42/233 (18.0)	203/512	<0.001
>3 wet nights/week	118/279 (42.3)	191/233 (82.0)	309/512	<0.001
<b>Daytime continence</b>				
Male	25/171 (14.6)	42/131 (32.1)	67/302	<0.001
Female	13/108 (12.0)	26/102 (25.5)	39/210	<0.001
Total	38/279 (13.6)	68/233 (29.2)	106/512	<0.001

Interestingly, most Chinese traditionally regarded faecal and/or urinary incontinence as normal for young infants and small children until the age of 4–5 years. Until recently, most people, and even medical practitioners, regarded bed-wetting as a normal phenomenon that does not warrant any intervention or treatment. Most parents would be reluctant to take a child with bed-wetting to seek medical advice until a very late age, and even if they do, the advice they receive from the medical practitioner most often would be simple observation, with a reassurance that the condition will disappear with time. This relative indifference or apathy within the Chinese community for PNE probably resulted in the low prevalence values reported in our previous epidemiological survey [6]. However, since the previous epidemiological study in the mid 1990s, an intensive public educational programme on NE has been introduced in Hong Kong, through a series of health talks, seminars, public lectures, featured articles, but most importantly via various mass media channels, including television and radio interviews, newspapers and magazines. In parallel with these there was a rapid increase in public awareness of PNE and the number of enuretic patients encountered in our clinic increased rapidly. Results from the present survey also show that more realistic epidemiological values can be obtained when the target study population has sufficient knowledge and interest in the problem, once misconceptions arising from traditional belief have been corrected.

The most important finding of the present study is that there were significant differences in characteristics between younger enuretic and older children. Although the overall prevalence of PNE decreased with increasing age, the proportion of patients with severe enuretic symptoms (wetting >3 nights/week) progressively increased. A significant proportion of adolescents (82%) had either moderate or severe enuresis, whereas most enuretic children (57.7%) have much milder bed-wetting with less than one enuretic episode/week. At age 5 years, only 14.3% of enuretic children wet 7 nights/week, compared with 48.3% at 19 years old ( $P < 0.001$ ). Notably, the results of a previous epidemiological study of PNE in adolescents and adults up to the age of 40 years indicated that the prevalence of PNE remained rather static, with no further significant decrease after the age of 10 years, and >2% of both

men and women remained enuretic. Of these affected adults, over half wet  $\geq 3$  nights/week, and a quarter had enuretic symptoms every night [10]. These findings therefore strongly suggest that the enuretic children with more severe symptoms probably have a significantly greater chance of persistent PNE in adult life. It is arguable therefore that for these groups of enuretic children with very severe symptoms, investigations and active treatments should be started at a much earlier age.

PNE is a heterogeneous disorder with various underlying pathophysiological mechanisms, causing in common a mismatch between the nocturnal bladder capacity and the amount of urine produced during sleep at night, in association with a simultaneous failure of conscious arousal, or waking, in response to the sensation of bladder fullness. Recent studies showed the important role of bladder dysfunction, e.g. small functional bladder capacity, instability during sleep, and detrusor hypercontractility caused by BOO, in the pathogenesis of PNE in children, especially those who are refractory to treatment [19–23]. Our previous epidemiological study of PNE in adolescents and adults showed a significantly higher incidence of urinary symptoms (frequency, urgency and incontinence), which were suggestive of underlying bladder dysfunction in adults with PNE, compared to normal controls [10]. Moreover, previous studies on bladder function in adult patients with PNE showed that the great majority (>90%) of adult enuretics had underlying detrusor overactivity, and 70% of the patients had urodynamic evidence of functional BOO [24]. This present study also showed that the incidence of daytime urinary symptoms was significantly higher in adolescents than younger enuretic children aged  $\leq 10$  years. The present finding suggesting that PNE will spontaneously resolve with age probably applies only to those with mild enuretic symptoms. Those with severe enuretic symptoms probably represent a more pronounced and refractory form of the condition, often associated with underlying bladder dysfunction, and would be more likely to have persistent enuretic symptoms into adult life [25–29]. This important finding may offer new clues to the poor treatment response in some enuretic patients with very severe symptoms. Further study of this group of adults with PNE, who in general have more pronounced and persistent symptoms than

younger enuretic children, may provide further insights into the complex pathophysiology of these various subtypes, and ways to refine the management of this heterogeneous disorder.

In conclusion, PNE is as prevalent in Hong Kong Chinese schoolchildren as in western populations. The present finding suggesting that PNE will spontaneously resolve with age probably applies only to those with mild enuretic symptoms; with increasing age there is a greater proportion of enuretic children with more severe bed-wetting. Enuretic children aged >10 years, and adolescents, have significantly more daytime urinary symptoms and incontinence. The previously reported low prevalence of PNE in Hong Kong was probably due to parental indifference to the problem.

#### ACKNOWLEDGEMENTS

The Health Care and Promotion Fund supported this project.

#### CONFLICT OF INTEREST

None declared.

#### REFERENCES

- 1 de Jonge GA. Epidemiology of enuresis: a survey of the literature. In Kovin I, MacKeith RC, Meadows SR eds, *Bladder Control and Enuresis*. London: Heinemann, 1973: 39–46
- 2 Fergusson DM, Hons BA, Horwood LJ, Shannon FT. Factors related to the age of attainment of nocturnal enuresis. *Behav Psychother* 1986; **78**: 884–90
- 3 Foxman B, Valdez RB, Brook RH. Childhood enuresis: prevalence, perceived impact, and prescribed treatments. *Pediatrics* 1986; **77**: 482–7
- 4 Hellstrom AL, Hansson E, Hansson S, Hjalmas K, Jodal U. Micturition habits and incontinence in 7-year-old Swedish school entrants. *Eur J Pediatr* 1990; **149**: 434–7
- 5 Watanabe H, Kawachi A. Nocturnal enuresis: social aspects and treatment perspectives in Japan. *Scan J Urol Nephrol Suppl* 1994; **163**: 29–38
- 6 Yeung CK. Nocturnal enuresis in Hong Kong: different Chinese phenotypes?

- Scan J Urol Nephrol Suppl* 1997; **183**: 17–21
- 7 **Cushing FC Jr, Baller WR.** The problem of nocturnal enuresis in adults: special reference to managers and managerial aspirants. *J Psychol* 1975; **89**: 203–13
  - 8 **van Son MJ, Mulder G, van Londen A.** The effectiveness of dry bed training for nocturnal enuresis in adults. *Behav Res Ther* 1990; **28**: 347–9
  - 9 **Tietjen DN, Husmann DA.** Nocturnal enuresis: a guide to evaluation and treatment. *Mayo Clin Proc* 1996; **71**: 857–62
  - 10 **Yeung CK, Sihoe JD, Sit FK, Bower W, Sreedhar B, Lau J.** Characteristics of primary nocturnal enuresis in adults: an epidemiological study. *BJU Int* 2004; **93**: 341–5
  - 11 **Wadsworth ML.** Persistent enuresis in adults. *Am J Orthopsychiatry* 1944; **14**: 313
  - 12 **Turner RK, Taylor PD.** Conditioning treatment of nocturnal enuresis in adults: preliminary findings. *Behav Res Ther* 1974; **12**: 41–52
  - 13 **Jarvelin MR, Vikevainen–Tervonen L, Moilanen I, Huttunen NP.** Enuresis in seven-year-old children. *Acta Paediatr Scand* 1988; **77**: 148–53
  - 14 **Mattson S.** Urinary incontinence and nocturia in healthy school children. *Acta Paediatr* 1944; **83**: 950–4
  - 15 **Bower WF, Moore KH, Shepherd R, Adams RD.** The epidemiological of childhood enuresis in Australia. *Br J Urol* 1996; **78**: 602–6
  - 16 **Byrd RS, Weitzman M, Lanphear NE, Auinger P.** Bedwetting in US children: epidemiology and related behaviour problems. *Pediatrics* 1996; **98**: 414–9
  - 17 **Serel TA, Akhan G, Koyuncuoglu R et al.** Epidemiology of enuresis in Turkish children. *Scand J Urol Nephrol* 1997; **31**: 537–9
  - 18 **Lee SD, Sohn DW, Lee JZ, Park NC, Chung MK.** An epidemiological study of enuresis in Korean children. *BJU Int* 2000; **85**: 869–73
  - 19 **Starfield B.** Functional bladder capacity in enuretic and nonenuretic children. *J Pediatr* 1967; **70**: 77–81
  - 20 **Eller DA, Homsy YL, Austin PF, Tanguay S, Cantor A.** Spot urine osmolality, age and bladder capacity as predictors of response to desmopressin in nocturnal enuresis. *Scand J Urol Nephrol Suppl* 1997; **183**: 41–5
  - 21 **Oredsson AF, Jorgensen TM.** Changes in nocturnal bladder capacity during the treatment with bell and pad for monosymptomatic nocturnal enuresis. *J Urol* 1998; **160**: 166–9
  - 22 **Yeung CK, Chiu HN, Sit FK.** Bladder dysfunction in children with refractory monosymptomatic primary nocturnal enuresis. *J Urol* 1999; **162**: 1049–55
  - 23 **Yeung CK, Sit FK, To LK et al.** Reduction in nocturnal functional bladder capacity is a common factor in the pathogenesis of refractory nocturnal enuresis. *BJU Int* 2002; **90**: 302–7
  - 24 **Yeung CK, Sihoe JD, Sit FK, Diao M, Yew SY.** Urodynamic findings in adults with primary nocturnal enuresis. *J Urol* 2004; **171**: 2595–8
  - 25 **Torrrens MJ, Collins CD.** The urodynamic assessment of adults enuresis. *Br J Urol* 1975; **47**: 433–40
  - 26 **Hirasing RA, van Leerdam FJ, Bolk–Bennink L, Janknegt RA.** Enuresis nocturna in adults. *Scand J Urol Nephrol* 1997; **31**: 533–6
  - 27 **Robertson G, Rittig S, Kovacs L, Gaskill MB, Zee P, Nanninga J.** Pathophysiology and treatment of enuresis in adults. *Scand J Urol Nephrol Suppl* 1999; **202**: 36–9
  - 28 **Burgio KL, Locher JL, Ives DG, Hardin JM, Newman AB, Kuller LH.** Nocturnal enuresis in community-dwelling older adults. *J Am Geriatr Soc* 1996; **44**: 139–43
  - 29 **Vandersteen DR, Husmann DA.** Treatment of primary nocturnal enuresis persisting into adulthood. *J Urol* 1999; **161**: 90–2

**Correspondence:** Chung K. Yeung, Chief, Division of Paediatric Surgery and Paediatric Urology, Department of Surgery, Chinese University of Hong Kong, Prince of Wales Hospital, Shatin, NT, Hong Kong, SAR, China. e-mail: ckyeung@surgery.cuhk.edu.hk

**Abbreviations:** PNE, primary nocturnal enuresis.